

# Combinatorial Principles and $\aleph_k$ -free Modules

Dissertation  
zur Erlangung des akademischen Grades eines  
Doktors der Naturwissenschaften

von

**Héctor Gabriel Salazar Pedroza**

vorgelegt bei der Fakultät für Mathematik  
der Universität Duisburg-Essen, Standort Essen

Essen 2012

**Antragssteller:** Héctor Gabriel Salazar Pedroza,  
geb. am 24.03.1982  
in León, Mexiko

**Abgabe der Dissertation:** 08.10.2012

**Zulassung zur Promotion:** 10.10.2012

**Tag der Disputation:** 26.11.2012

**Prüfungsausschuss:**

Vorsitzender:	Prof. Dr. Karl-Josef Witsch
Gutachter:	Prof. Dr. Ulrich Albrecht
	Prof. Dr. Rüdiger Göbel

*A mi esposa Marcela.*

## Contents

Introduction	v
Acknowledgements	ix
List of Symbols	x
Chapter 1. Separable $\aleph_k$ -free Groups with Small Dual	1
1. Set-theoretic Preliminaries	1
2. The Construction	10
3. Slenderness	17
Chapter 2. Separable $\aleph_k$ -free Groups Satisfying $\text{End } G = A \oplus \text{Fin } G$	21
1. Set-theoretic Modifications	21
2. Triple Modules	26
3. The Step Lemma	39
4. The Final Construction	51
References	59

## Introduction

As part of his PhD-thesis at Cambridge University, A.L.S. Corner obtained the following

**THEOREM 1.** *Every countable reduced torsion-free ring  $R$  with  $1$  is isomorphic to the endomorphism ring  $\text{End } G$  of a countable reduced torsion-free group  $G$ .*

This powerful theorem (in [1]) is very important since it permits solving various problems by choosing different rings for  $R$ . For example, this allows to give a negative answer to Kaplansky's Test Problems. Years later this result was extended by A.L.S. Corner and R. Göbel in [2] to modules over rings of uncountable size using a combinatorial principle by S. Shelah called the "Black Box". This also allows to construct  $E$ -rings which are now widely used in algebraic topology. It was already noted in [2] that the constructed modules were  $\aleph_1$ -free (i.e. every  $< \aleph_1$ -generated submodule is contained in a free submodule) but it was clear that they were not  $\aleph_2$ -free. Shortly after another paper by M. Dugas and R. Göbel [5] discussed separable groups, in which every finite subset is contained in a free direct summand of the group. In the case of torsion-free abelian groups this means that the group is a pure subgroup of a Cartesian product of copies of the integers. The main result in this paper can be summarized in the following

**THEOREM 2.** *For every infinite cardinal  $\lambda$  such that  $\lambda^{\aleph_0} = \lambda$  there exists a separable  $\aleph_1$ -free abelian group  $G$  with  $\text{End } G = \mathbb{Z} \oplus \text{Fin } G$ , where  $\text{Fin } G$  is the ideal generated by all the endomorphisms of  $G$  whose images have finite rank.*

Already in the first half of the last century Baer, Specker ([10], vol. 1, p. 94) and Łoś ([10], vol. 2, p. 161) observed that there are complicated  $\aleph_1$ -free groups that are not free. One famous example of such groups is the Baer-Specker group  $\prod_{n < \omega} \mathbb{Z}e_n$ . G. Nöbeling and G.M. Bergman ([10], vol. 2, p. 173, 174) observed that these groups contain large “canonical” free subgroups. It is natural to ask if we can strengthen the theorem from [2] to obtain  $\aleph_k$ -free modules, namely, can we find complicated  $\aleph_k$ -free modules which are not free for  $k > 1$ ? Classical work in this direction was done by P. Griffith [16], P. Hill [17], P. Eklof and S. Shelah [9]. While this can be relatively easily done assuming additional set-theoretic assumptions such as the axiom of constructibility  $V = L$  (see Dugas, Göbel [4]), the construction of such modules within ZFC without extra axioms is a complicated target.

Göbel and Shelah [12] follow this approach by using a modification of the Black Box in their proof:

**THEOREM 3.** *For every  $k < \omega$  there exists an  $\aleph_k$ -free abelian group  $G$  of size  $\beth_k$  such that  $\text{Hom}(G, \mathbb{Z}) = 0$ .*

In 2010 R. Göbel, D. Herden and S. Shelah [11] developed a new, more powerful version of the Black Box principle, called the  $\aleph_k$ -Black Box, which allowed them to prove the following

THEOREM 4. *Let  $R$  be a  $p$ -cotorsion-free domain and  $A$  an  $R$ -algebra with free  $R$ -module  $A_R$  and  $|A| \leq \mu$ . If  $\lambda = \beth_k^+(\mu)$  for some positive integer  $k$ , where  $\beth_0^+(\mu) = \mu$  and  $\beth_{n+1}^+(\mu) = \left(2^{\beth_n^+(\mu)}\right)^+$ , which is the successor cardinal of the powerset of  $\beth_n^+(\mu)$ , then we can construct an  $\aleph_k$ -free  $A$ -module  $G$  of cardinality  $\lambda$  with  $R$ -endomorphism algebra  $\text{End}_R G = A$ .*

The two main results of this work deal with the construction of separable  $\aleph_k$ -free abelian groups following the approach of Theorem 3 and Theorem 4. It is therefore natural to keep the notation as similar as possible to the ones used in these papers. We cannot expect these groups to have a trivial dual  $\text{Hom}(G, \mathbb{Z})$  or satisfy  $\text{End } G = A$ , because separable groups embed into cartesian products  $\prod_{i \in I} \mathbb{Z}e_i$ . The projections induced by these products cannot be eliminated and require to consider the ideal  $\text{Fin } G$ .

The first result of this work is the construction of a separable  $\aleph_k$ -free abelian group  $G$  of size  $\beth_k$  with no epimorphisms onto  $\bigoplus_{n < \omega} \mathbb{Z}e_n$ , and is based on methods used in the proof of Theorem 3:

THEOREM 5. *For every  $k > 1$  there exists a separable  $\aleph_k$ -free abelian group  $G$  of size  $\beth_k$  with no epimorphisms onto  $\bigoplus_{n < \omega} \mathbb{Z}e_n$ .*

The second result shows the existence of an  $\aleph_k$ -free  $A$ -module  $G$  of size  $\beth_k^+(\mu)$  which is separable as an abelian group and satisfies  $\text{End } G = A \oplus \text{Fin } G$ , where  $A$  is a given ring ( $\mathbb{Z}$ -algebra) with  $|A| \leq \mu$  and free additive structure  $A^+ = \bigoplus_{\alpha < \kappa} \mathbb{Z}e_\alpha$ . Here we follow [11]:

**THEOREM 6.** *For every  $k > 1$  there exist arbitrarily large  $\aleph_k$ -free  $A$ -modules  $G$ , which are separable as abelian groups, such that  $\text{End } G = A \oplus \text{Fin } G$ .*

This work is presented in two chapters, one for each of these main results. The first section of Chapter 1 introduces the basic notations and definitions, in particular the definition of the basic sets  $\Lambda$  and  $\Lambda_*$ . We also mention the set-theoretic version of the new Black Box. Section 2 presents the new Black Box in an algebraic context and constructs separable  $\aleph_k$ -free groups  $G$  satisfying  $\text{Hom}(G, H) = \text{Fin}(G, H)$  for  $H$  a free abelian group. In Section 3 we briefly discuss the slenderness of  $\aleph_k$ -free groups.

The first section of Chapter 2 considers a ring  $A$  with free additive structure. After some slight modifications to the set-theoretic notions and definitions of Chapter 1, we construct  $\aleph_k$ -free  $A$ -modules, which are separable abelian groups, by means of pairs of subsets of our basic sets. Section 2 introduces another kind of  $A$ -modules with important freeness properties, which are defined by means of triples of subsets of these basic sets. The focus of Section 3 is the Step Lemma, which is the central piece of the final construction. Finally, Section 4 presents the main construction with the help of the Strong Black Box and a result based on [5].



## Acknowledgements

I am deeply grateful to my supervisor Rüdiger Göbel for all the dedication, advice and support he offered me for the realization of this work during my stay in Essen.

I would like to thank Ulrich Albrecht for all the valuable comments he made to improve this work.

I would like to express my great appreciation to all the current and former members of the AG Göbel I met in Essen for a pleasant working environment. I am specially in dept to Daniel Herden for all the time he dedicated to helping me inside and outside our working place.

I thank my wife Marcela dearly for all her love, understanding and patience. Also many thanks to my parents, siblings and friends for their invaluable help and support.

Last but not least, I want to thank the DAAD for awarding me with the PhD-grant that made this work and this great experience possible. <sup>1</sup>

---

<sup>1</sup>This work was supported by the Deutscher Akademischer Austauschdienst (German Academic Exchange Service) by means of a “DAAD-Forschungsstipendium für Doktoranden und Nachwuchswissenschaftler”.

## List of Symbols

$\beth_n^+(\mu)$ , vii

$\omega\lambda$ ,  $\omega^\uparrow\lambda$ ,  $\omega^{>}\lambda$ ,  $\omega^{\uparrow>}\lambda$ , 1

$[A]^{\leq\kappa}$ ,  $[A]^{<\kappa}$ ,  $[A]^\kappa$ , 1

$[\alpha, \gamma]$ ,  $(\alpha, \gamma)$ ,  $[\alpha, \gamma)$ , 1

$a \wedge b$ ,  $X \wedge Y$ , 2

$\Lambda^{[i,j]}$ ,  $\Lambda_m^{[i,j]}$ ,  $\Lambda_*^{[i,j]}$ ,  $\Lambda$ ,  $\Lambda_*$ , 2

$[\eta]$ , 2

$\bar{\eta} \restriction \langle m, n \rangle$ , 3

$[\bar{\eta} \restriction m]_n$ ,  $[\bar{\eta} \restriction m]$ , 3

$[\bar{\eta}]_n$ ,  $[\bar{\eta}]$ , 3

$\|X\|$ , 3

$\bar{\lambda}$ , 8

$B_{Y_*}$ , 10, 21

$B$ ,  $\overline{B}$ , 10, 22

$[b]$ , 10

$y_{\bar{\eta}m}$ ,  $y_{\bar{\eta}}$ , 11

$b_i^{\bar{\eta}}$ ,  $b_{\bar{\eta}n}$ ,  $b_{\bar{\eta}}$ , 11

$y'_{\bar{\eta}m}$ ,  $y'_{\bar{\eta}}$ , 11

$[g]_\Lambda$ ,  $[H]_\Lambda$ , 13

$\overline{A}$ , 21

$\mathfrak{F}_{Y_*Y}$ ,  $\mathfrak{F}_{Y_*X}$ , 22

$G_{Y_*Y}$ , 23

$Y_{X_*}$ , 26

$u_{\bar{\eta}}(X_*)$ , 26

$\rho_{Y_*X_*}$ , 26

$G_{Y_*YX_*}$ , 26

$Y(X_*)$ , 30

$\text{Cp}(X_*)$ , 35

$\text{PC}(X_*, Y_*)$ , 35

$\Lambda^{\bar{\xi}}$ ,  $\Lambda_*^{\bar{\xi}}$ ,  $\Lambda^{\bar{\xi}*}$ , 39

$\Lambda_{[\eta[n]]}^{\bar{\xi}}$ ,  $\Lambda_{\eta}^{\bar{\xi}*}$ , 39

## CHAPTER 1

### Separable $\aleph_k$ -free Groups with Small Dual

#### 1. Set-theoretic Preliminaries

Let us begin by introducing some notation:

- (1) Functions will be written on the right side of their argument, so if  $f$  is a function with domain  $A$  and  $a \in A$ , then the image of  $a$  under  $f$  will be written as  $af$ .
- (2)  ${}^\omega\lambda$  denotes the set of all functions  $\tau : \omega \rightarrow \lambda$ , while  ${}^{\omega\uparrow}\lambda$  is the subset of  ${}^\omega\lambda$  consisting of all *order preserving* functions  $\eta : \omega \rightarrow \lambda$ , namely

$${}^{\omega\uparrow}\lambda = \{ \eta : \omega \rightarrow \lambda \mid m\eta < n\eta \text{ for } m < n \}.$$

Similarly,  ${}^{\omega>}\lambda$  denotes the set of all functions  $\sigma : n \rightarrow \lambda$  with  $n < \omega$ , while  ${}^{\omega\uparrow>}\lambda$  is the subset of  ${}^{\omega>}\lambda$  consisting of all *order preserving* functions  $\eta : n \rightarrow \lambda$  with  $n < \omega$ .

- (3) If  $f : A \rightarrow {}^BC$ , i.e.  $af$  is a function for all  $a \in A$ , then we write  $f_a$  instead of  $af$ .
- (4) If  $A$  is a set and  $\kappa$  is a cardinal, then  $[A]^{\leq\kappa}$  denotes the set of all  $X \subseteq A$  such that  $|X| \leq \kappa$ . Analogously we define  $[A]^{<\kappa}$  and  $[A]^\kappa = [A]^{=\kappa}$ .
- (5) If  $\alpha \leq \gamma$  are ordinals, we write  $[\alpha, \gamma] = \{ \beta \mid \alpha \leq \beta \leq \gamma \}$ ,  $(\alpha, \gamma) = \{ \beta \mid \alpha < \beta < \gamma \}$  and  $[\alpha, \gamma) = \{ \beta \mid \alpha \leq \beta < \gamma \}$ .
- (6) The expression  $H \leq G$  denotes that  $H$  is a subgroup of  $G$ .

- (7) Let  $\{A_i \mid i \in [1, m]\}$  and  $\{B_i \mid i \in [1, n]\}$  be finite families of sets,  $A = A_1 \times \cdots \times A_m$  and  $B = B_1 \times \cdots \times B_n$ . If  $a = (a_1, \dots, a_m) \in A$  and  $b = (b_1, \dots, b_n) \in B$ , we write  $a \wedge b = (a_1, \dots, a_m, b_1, \dots, b_n)$ . If  $X \subseteq A$  and  $Y \subseteq B$ , we write  $X \wedge Y = \{a \wedge b \mid a \in X, b \in Y\}$ .

Let  $k > 1$  be fixed for the rest of this work. Given a finite sequence of infinite cardinals  $\langle \lambda_1, \dots, \lambda_k \rangle$  and  $i, j \in [1, k]$  with  $i \leq j$ , we construct the set

$$\Lambda^{[i,j]} = \omega^\uparrow \lambda_i \times \omega^\uparrow \lambda_{i+1} \times \cdots \times \omega^\uparrow \lambda_{j-1} \times \omega^\uparrow \lambda_j.$$

For  $\Lambda^{[1,k]}$  we simply write  $\Lambda$ . Moreover, for  $i < j$ , define  $\Lambda_i^{[i,j]} = \omega^{\uparrow >} \lambda_i \wedge \Lambda^{[i+1,j]}$ ,  $\Lambda_j^{[i,j]} = \Lambda^{[i,j-1]} \wedge \omega^{\uparrow >} \lambda_j$  and

$$\Lambda_m^{[i,j]} = \Lambda^{[i,m-1]} \wedge \omega^{\uparrow >} \lambda_m \wedge \Lambda^{[m+1,j]},$$

for all  $m \in (i, j)$ . Put

$$\Lambda_*^{[i,j]} = \bigcup_{m \in [i,j]} \Lambda_m^{[i,j]}.$$

For  $i = 1$  and  $j = k$  we simply write  $\Lambda_*$  and  $\Lambda_m$  for all  $m \in [1, k]$ .

DEFINITION 7. (1) If  $\eta \in \omega^\uparrow \lambda_m$ , then the *support* of  $\eta$  is the set

$$[\eta] = \{\eta \upharpoonright n \mid n < \omega\}.$$

(2) If  $\nu \in \omega^{>\uparrow} \lambda_m$  with  $\text{dom } \nu = n$ , then the *support* of  $\nu$  is the set

$$[\nu] = \{\nu \upharpoonright \ell \mid \ell \leq n\}.$$

- (3) If  $\bar{\eta} = (\eta_i, \dots, \eta_j) \in \Lambda^{[i,j]}$ ,  $m \in [i, j]$  and  $n < \omega$ , then  $\bar{\eta} \restriction \langle m, n \rangle$  denotes the element of  $\Lambda_m^{[i,j]}$  obtained from  $\bar{\eta}$  by replacing its component  $\eta_m$  by  $\eta_m \restriction n$ , i.e.

$$\bar{\eta} \restriction \langle m, n \rangle = (\eta_i, \dots, \eta_{m-1}, \eta_m \restriction n, \eta_{m+1}, \dots, \eta_j).$$

- (4) For every  $\bar{\eta} \in \Lambda^{[i,j]}$ ,  $m \in [i, j]$  and  $n < \omega$  consider the sets

$$[\bar{\eta} \restriction m]_n = \{ \bar{\eta} \restriction \langle m, n' \rangle \mid n' \in [n, \omega) \}$$

and

$$[\bar{\eta}]_n = \bigcup_{m \in [i,j]} [\bar{\eta} \restriction m]_n$$

If  $n = 0$ , then we simply drop the subindex and write  $[\bar{\eta} \restriction m]$  and  $[\bar{\eta}]$  instead.

The set  $[\bar{\eta}]$  is called the *support* of  $\bar{\eta}$ .

DEFINITION 8. Define the *norm function*  $\|\cdot\| : \Lambda \dot{\cup} \Lambda_* \rightarrow \lambda_k$  as

$$\|\bar{\eta}\| = \sup \text{Im } \eta_k.$$

For  $X \subseteq \Lambda \dot{\cup} \Lambda_*$ , we naturally define

$$\|X\| = \sup_{\bar{\eta} \in X} \|\bar{\eta}\|.$$

DEFINITION 9. A function  $F : \Lambda \rightarrow [\Lambda_*]^{\leq \aleph_0}$  is *regressive* if  $\|\bar{\eta}F\| < 0\eta_k$  for all  $\bar{\eta} \in \Lambda$ .

The first step towards the proof of the main result concerning  $\aleph_k$ -freeness is the so called Freeness-Proposition, which allows us to enumerate subsets of  $\Lambda$  in such a “clever” way that we can prove linearly independence in the constructed groups. First we have to deal with the notion of coordinatewise-closedness of filtrations, which will appear in the proof of the proposition.

LEMMA 10. *Let  $F : \Lambda \rightarrow [\Lambda_*]^{\leq \aleph_0}$  be any map,  $f \in [1, k]$  and  $\Omega \in [\Lambda]^{\aleph_f}$  together with an  $\aleph_f$ -filtration  $\{\mathcal{U}^\alpha \mid \alpha < \omega_f\}$  of  $\Omega$  such that  $\mathcal{U}^0 = \emptyset$  and  $|\mathcal{U}^{\alpha+1} \setminus \mathcal{U}^\alpha| = \aleph_{f-1}$  for all  $\alpha < \omega_f$ . Then it is possible to construct a coordinatewise-closed  $\aleph_f$ -filtration  $\{\Omega^\alpha \mid \alpha < \omega_f\}$  of  $\Omega$ , meaning that for all  $\bar{\eta} \in \Omega^{\alpha+1}$ , if there exist  $\bar{\eta}', \bar{\eta}'' \in \Omega^\alpha$  such that*

$$\{\eta_m \mid m \in [1, k]\} \subseteq \{\eta'_m, \eta''_m \mid m \in [1, k]\} \cup \{\nu_m \mid \bar{\nu} \in \bar{\eta}' F \cup \bar{\eta}'' F, m \in [1, k]\},$$

*then  $\bar{\eta} \in \Omega_\alpha$ . This  $\aleph_f$ -filtration also satisfies  $\Omega^0 = \emptyset$  and  $|\Omega^{\alpha+1} \setminus \Omega^\alpha| = \aleph_{f-1}$  for all  $\alpha < \omega_f$ .*

PROOF. First suppose that we have constructed a coordinatewise-closed  $\aleph_f$ -filtration  $\{\Omega^\alpha \mid \alpha < \gamma\}$  of  $\Omega$  up to some limit ordinal  $\gamma < \omega_f$  such that  $\Omega^0 = \emptyset$  and  $|\Omega^{\alpha+1} \setminus \Omega^\alpha| = \aleph_{f-1}$  for all  $\alpha + 1 < \gamma$  by means of the original filtration. We define  $\Omega^\gamma = \bigcup_{\alpha < \gamma} \Omega^\alpha$  as usual.

Now suppose that the coordinatewise-closed  $\aleph_f$ -filtration  $\{\Omega^\alpha \mid \alpha \leq \gamma\}$  of  $\Omega$  has been constructed up to some successor ordinal  $\gamma < \omega_f$  and also satisfies  $|\Omega^{\alpha+1} \setminus \Omega^\alpha| = \aleph_{f-1}$  for all  $\alpha + 1 \leq \gamma$ . Let  $\beta$  be the minimal ordinal such that  $\Omega^\gamma \subseteq \mathcal{U}^\beta$  and  $|\mathcal{U}^\beta \setminus \Omega^\gamma| = \aleph_{f-1}$ . Let  $\Omega_0^{\gamma+1} = \mathcal{U}^\beta$  and assume we have constructed  $\Omega_n^{\gamma+1}$  for some  $n < \omega$ . For all

$m \in [1, k]$ , let

$$\omega^\uparrow \lambda_m(\Omega_n^{\gamma+1}) = \left\{ \eta_\ell \mid \ell \in [1, k], \bar{\eta} \in \Omega_n^{\gamma+1} \cup \bigcup \Omega_n^{\gamma+1} F \right\} \cap \omega^\uparrow \lambda_m,$$

$$\Omega_{n+1}^{\gamma+1} = \omega^\uparrow \lambda_1(\Omega_n^{\gamma+1}) \times \cdots \times \omega^\uparrow \lambda_k(\Omega_n^{\gamma+1}) \subseteq \Lambda$$

and

$$\Omega^{\gamma+1} = \Omega \cap \bigcup_{n < \omega} \Omega_n^{\gamma+1}.$$

□

**FREENESS-PROPOSITION 11.** *Let  $F : \Lambda \rightarrow [\Lambda_*]^{\leq \aleph_0}$  be a regressive map,  $f \in [1, k]$ ,  $\Omega \in [\Lambda]^{\aleph_{f-1}}$  and  $\langle u_{\bar{\eta}} \mid \bar{\eta} \in \Omega \rangle$  be a family of subsets of  $[1, k]$  such that  $|u_{\bar{\eta}}| \geq f$ . Then there exists a bijective enumeration  $\{\bar{\eta}^\alpha \mid \alpha < \zeta\}$  of  $\Omega$  for some  $\zeta \in [\omega_{f-1}, \omega_f)$  such that, for all  $\alpha < \zeta$ , there exist  $\ell_\alpha \in u_{\bar{\eta}^\alpha}$  and  $n_\alpha \in [1, \omega)$  with the property that, for all  $n \geq n_\alpha$ ,*

$$\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle \notin \{ \bar{\eta}^\beta \restriction \langle \ell_\alpha, n \rangle \mid \beta < \alpha \} \cup \bigcup \Omega_\alpha F$$

where  $\Omega_\alpha = \{ \bar{\eta}^\beta \mid \beta \leq \alpha \}$ .

**PROOF.** We proceed by induction on  $f$ . If  $f = 1$ , then  $|\Omega| = \aleph_0$  and  $u_{\bar{\eta}} \neq \emptyset$  for all  $\bar{\eta} \in \Omega$ . For all  $\alpha < \aleph_k$ , define  $U_\alpha = \{ \bar{\eta} \in \Omega \mid 0\eta_k = \alpha \}$ . Let  $N = \{ \alpha < \aleph_k \mid U_\alpha \neq \emptyset \}$  and enumerate it  $N = \{ \alpha_\beta \mid \beta < \delta \}$  for some  $\delta < \omega_1$  in such a way that  $\alpha_\beta < \alpha_\gamma$  if and only if  $\beta < \gamma < \delta$ . Put  $\gamma_\beta = |U_{\alpha_\beta}|$  and  $\sigma_\beta = \sum_{\alpha < \beta} \gamma_\alpha$ . We enumerate  $U_{\alpha_\beta} = \{ \bar{\eta}^\alpha \mid \sigma_\beta \leq \alpha < \sigma_\beta + \gamma_\beta \}$ . This results in a bijective enumeration  $\{ \bar{\eta}^\alpha \mid \alpha < \zeta \}$  of  $\Omega$  such that  $\zeta \in [\omega, \omega_1)$  and, for all  $\alpha < \zeta$ ,

$$0\eta_k^\alpha \leq 0\eta_k^{\alpha+1} < 0\eta_k^{\alpha+\omega}.$$

Choose  $\ell_\alpha \in u_{\bar{\eta}^\alpha}$  arbitrarily. If  $\bar{\eta}^\alpha \in U_\gamma$  and  $\beta_0$  is the minimal ordinal such that  $\bar{\eta}^{\beta_0} \in U_\gamma$ , then we can find some  $n_{\alpha,\beta} \in [1, \omega)$  such that  $\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle \neq \bar{\eta}^\beta \restriction \langle \ell_\alpha, n \rangle$  for all  $\beta \in [\beta_0, \alpha)$  and  $n \geq n_{\alpha,\beta}$ . Put  $n_\alpha = \max_{\beta \in [\beta_0, \alpha)} n_{\alpha,\beta}$ . Then, for all  $n \geq n_\alpha$ ,  $\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle \notin \{ \bar{\eta}^\beta \restriction \langle \ell_\alpha, n \rangle \mid \beta \in [\beta_0, \alpha) \}$ . Moreover,

$$\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle \notin \{ \bar{\eta}^\beta \restriction \langle \ell_\alpha, n \rangle \mid \beta < \beta_0 \} \cup \bigcup \Omega_\alpha F$$

since  $F$  is regressive and  $0\eta_k^\beta < 0\eta_k^\alpha$  for all  $\beta < \beta_0$ .

Now suppose that the assertion is true for some  $f \in [1, k-1]$ . Let  $\Omega \in [\Lambda]^{\aleph_f}$  and  $\langle u_{\bar{\eta}} \mid \bar{\eta} \in \Omega \rangle$  with  $|u_{\bar{\eta}}| \geq f+1$ . Choose an  $\aleph_f$ -filtration  $\{ \Omega^\alpha \mid \alpha < \omega_f \}$  of  $\Omega$  such that  $\Omega^0 = \emptyset$  and  $|\Omega^{\alpha+1} \setminus \Omega^\alpha| = \aleph_{f-1}$  for all  $\alpha < \omega_f$ . By the previous lemma, we can assume that this filtration is coordinatewise-closed. For every  $\bar{\eta} \in \Omega^{\alpha+1} \setminus \Omega^\alpha$ , consider

$$u_{\bar{\eta}}^* = \{ m \in [1, k] \mid \exists \bar{\eta}' \in \Omega^\alpha, n < \omega (\bar{\eta} \restriction \langle m, n \rangle = \bar{\eta}' \restriction \langle m, n \rangle \text{ or } \bar{\eta} \restriction \langle m, n \rangle \in \bar{\eta}' F) \}.$$

It follows that  $|u_{\bar{\eta}}^*| \leq 1$ , since  $|u_{\bar{\eta}}^*| > 1$  would imply that  $\bar{\eta} \in \Omega^\alpha$ . Put  $u'_{\bar{\eta}} = u_{\bar{\eta}} \setminus u_{\bar{\eta}}^*$  and observe that  $|u'_{\bar{\eta}}| \geq f$ . We apply the induction hypothesis on each of the sets  $\Omega^{\alpha+1} \setminus \Omega^\alpha$  together with the family  $\langle u'_{\bar{\eta}} \mid \bar{\eta} \in \Omega^{\alpha+1} \setminus \Omega^\alpha \rangle$  to obtain an enumeration  $\Omega^{\alpha+1} \setminus \Omega^\alpha = \langle \bar{\eta}^\beta \mid \beta < \zeta \rangle$  for some  $\zeta \in [\omega_{f-1}, \omega_f)$  with the required property. We induce an enumeration on  $\Omega$  with the desired property by ordering these enumerations lexicographically.  $\square$

Now we get ready to present the  $\bar{\lambda}$ -Black Box, the prediction principle we will use to get rid of unwanted homomorphisms. Please keep our notation (3) from page 1 in mind.



LEMMA 12. *Let  $\lambda$  be an infinite cardinal and  $\mathfrak{P}$  a set of cardinality  $\leq \lambda^{\aleph_0}$ . Then there exists a map  $\Phi : {}^\omega\lambda \rightarrow {}^\omega\mathfrak{P}$  such that, for all  $\mathfrak{f} : {}^{\omega^>}\lambda \rightarrow \mathfrak{P}$  and  $\nu \in {}^{\omega^>}\lambda$ , we can find  $\eta \in {}^\omega\lambda$  with  $\nu \subset \eta$  and  $n\Phi_\eta = (\eta \restriction n)\mathfrak{f}$  for all  $n < \omega$ .*

PROOF. Since  $|\mathfrak{P}| \leq \lambda^{\aleph_0} = |{}^\omega\lambda|$ , we can fix an embedding  $\pi : \mathfrak{P} \hookrightarrow {}^\omega\lambda$ . We also fix a map  $\mu : \lambda \rightarrow {}^{\omega^>}\lambda$  such that the preimage  $\mu^{-1}[\sigma]$  is unbounded in  $\lambda$  for all  $\sigma \in {}^{\omega^>}\lambda$ .

We would like to identify every function in  ${}^n\mathfrak{P}$  with a function in  ${}^{\omega^>}\lambda$  for all  $n < \omega$ . For this reason we define a *coding map*  $\pi^n : {}^n\mathfrak{P} \rightarrow {}^n\lambda$  for all  $n < \omega$  such that if  $\varphi \in {}^n\mathfrak{P}$ , then  $\pi_\varphi^n$  is given by  $(qn + r)\pi_\varphi^n = r\pi_{q\varphi}$  for  $q, r \in [0, n)$ . We now consider the set

$$X = \{ \delta \in {}^\omega\lambda \mid \exists \psi_\delta \in {}^\omega\mathfrak{P} \exists i_\delta < \omega \forall n \geq i_\delta (\pi_{\psi_\delta \restriction n}^n = \mu_{n\delta}) \}.$$

Because  $\pi$  is an embedding, we obtain that if  $\delta \in X$ , then  $\psi_\delta$  is unique. We use  $X$  to define  $\Phi$  in the following way: if  $\delta \notin X$ , we take an arbitrary  $\Phi_\delta \in {}^\omega\mathfrak{P}$ , and if  $\delta \in X$ , we define  $\Phi_\delta = \psi_\delta$ .

Let  $\mathfrak{f} : {}^{\omega^>}\lambda \rightarrow \mathfrak{P}$  and  $\rho \in {}^{\omega^>}\lambda$ . Since we need to show that there is an extension  $\eta \in {}^\omega\lambda$  of  $\rho$ , we define  $n\eta = n\rho$  for all  $n \in \text{dom } \rho$ . Now assume we have defined  $\eta \restriction n$  up to a certain  $n \geq \text{dom } \rho$ . Consider the element  $\psi_{\eta n} \in {}^n\mathfrak{P}$  given by  $m\psi_{\eta n} = (\eta \restriction m)\mathfrak{f}$  for all  $m < n$ . Then  $\pi_{\psi_{\eta n}}^n \in {}^n\lambda$  and  $\mu^{-1}[\pi_{\psi_{\eta n}}^n]$  is unbounded in  $\lambda$ . Define  $n\eta = \alpha$  to be the least ordinal  $\alpha > (n-1)\eta$  such that  $\mu_\alpha = \pi_{\psi_{\eta n}}^n$ . This finishes the construction of the extension  $\eta \in {}^\omega\lambda$  of  $\rho$ . Moreover, let  $\psi_\eta = \bigcup_{n \in [\text{dom } \rho, \omega)} \psi_{\eta n}$ . Since  $\eta \in X$  is witnessed by  $\psi_\eta$  and  $i_\eta = \text{dom } \rho$ , it immediately follows that  $n\Phi_\eta = n\psi_\eta = (\eta \restriction n)\mathfrak{f}$  for all  $n < \omega$ .  $\square$

DEFINITION 13. A finite sequence of cardinals  $\bar{\lambda} = \langle \lambda_1, \dots, \lambda_k \rangle$  is a  $\blacksquare$ -sequence (*Black Box sequence*) if for every  $m \in [1, k]$ , the cardinals  $\chi_m = \lambda_m^{\aleph_0}$  satisfy

$$\chi_{m+1}^{\chi_m} = \chi_{m+1}.$$

DEFINITION 14. Let  $\bar{C} = \langle C_1, \dots, C_k \rangle$  be a sequence of sets such that  $|C_m| \leq \chi_m$  and take  $C = \bigcup_{m \in [1, k]} C_m$ . A *set-trap* for  $\bar{\eta} \in \Lambda$  and  $\bar{C}$  is a function  $\varphi_{\bar{\eta}} : [\bar{\eta}] \rightarrow C$ .

THE  $\bar{\lambda}$ -BLACK BOX 15. Let  $\bar{\lambda} = \langle \lambda_1, \dots, \lambda_k \rangle$  be any  $\blacksquare$ -sequence,  $\Lambda$  and  $\Lambda_*$  as before,  $\bar{C} = \langle C_1, \dots, C_k \rangle$  and  $C$  as in Definition 14. Then there exists a family of set-traps  $\langle \varphi_{\bar{\eta}} \mid \bar{\eta} \in \Lambda \rangle$  satisfying the following

Prediction Principle: If  $\varphi : \Lambda_* \rightarrow C$  is any map with the trap condition  $\Lambda_m \varphi \subseteq C_m$  and  $\alpha < \lambda_k$ , then there exists  $\bar{\eta} \in \Lambda$  such that  $\varphi \upharpoonright [\bar{\eta}] = \varphi_{\bar{\eta}}$  and  $0\eta_k = \alpha$ .

PROOF. We will proceed by induction on the length of  $\bar{\lambda}$ .

Assume that the length of  $\bar{\lambda}$  is 1, so we can simply write  $\bar{\lambda} = \langle \lambda \rangle$  and  $\bar{C} = \langle C \rangle$ . Then  $\Lambda = {}^\omega \lambda$  and  $\Lambda_* = {}^{\omega \uparrow} \lambda$ . Since  $|C| \leq \chi = \lambda^{\aleph_0}$ , Lemma 12 provides us with a map  $\Phi : \Lambda \rightarrow {}^\omega C$ . For all  $\eta \in \Lambda$  and  $n < \omega$ , define  $(\eta \upharpoonright n)\varphi_\eta = n\Phi_\eta$  to obtain a family of set-traps  $\langle \varphi_\eta \mid \eta \in \Lambda \rangle$ . Now let  $\varphi : \Lambda_* \rightarrow C$  be a map (here the condition  $\Lambda_* \varphi \subseteq C$  is redundant), and suppose  $\alpha < \lambda$ . Choose an arbitrary  $\nu \in \Lambda_*$  such that  $0\nu = \alpha$ . By Lemma 12, there exists  $\eta \in \Lambda$  such that  $\nu \subset \eta$  and  $n\Phi_\eta = (\eta \upharpoonright n)\varphi$  for all  $n < \omega$ . This means  $\varphi \upharpoonright [\eta] = \varphi_\eta$  and  $0\eta = \alpha$ .

Now assume that the assertion is true for some  $f \in [1, k-1]$  and that the length of  $\bar{\lambda}$  is  $f+1$ . In this case,  $\bar{\lambda} = \langle \lambda_1, \dots, \lambda_{f+1} \rangle$  and  $\bar{C} = \langle C_1, \dots, C_{f+1} \rangle$ . We also

write  $C^m = \bigcup_{i=1}^m C_i$  for  $m \in [1, f+1]$ . Define  $\mathfrak{P} = {}^{\Lambda^{[1,f]}}C_{f+1}$  (the set of all maps from  $\Lambda^{[1,f]}$  to  $C_{f+1}$ ) and notice that  $|\mathfrak{P}| \leq \chi_{f+1}^{\chi_f} = \chi_{f+1} = \lambda_{f+1}^{\aleph_0}$ . Hence, Lemma 12 provides us with a map  $\Phi : {}^{\omega^\uparrow}\lambda_{f+1} \rightarrow {}^\omega\mathfrak{P}$ . We would like to define the set-traps  $\varphi_{\bar{\eta}}^{f+1}$  for every  $\bar{\eta} \in \Lambda^{[1,f+1]}$ . By induction hypothesis, we already have a family of set-traps  $\langle \varphi_{\bar{\eta}}^f \mid \bar{\eta} \in \Lambda^{[1,f]} \rangle$ . Given an  $\bar{\eta} = (\eta_1, \dots, \eta_{f+1}) \in \Lambda^{[1,f+1]}$ , put  $\bar{\eta}' = (\eta_1, \dots, \eta_f) \in \Lambda^{[1,f]}$ . Then for every  $\bar{\eta} \in \Lambda^{[1,f+1]}$  define

$$(\bar{\eta} \restriction \langle m, n \rangle) \varphi_{\bar{\eta}}^{f+1} = \begin{cases} (\bar{\eta}' \restriction \langle m, n \rangle) \varphi_{\bar{\eta}'}^f, & \text{if } m \in [1, f]; \\ \bar{\eta}' \Phi_{\eta_{f+1}n}, & \text{if } m = f+1. \end{cases}$$

It remains to verify the prediction principle. Let  $\varphi : \Lambda_*^{[1,f+1]} \rightarrow C^{f+1}$  be such that  $\Lambda_m^{[1,f+1]} \varphi \subseteq C_m$  and  $\alpha < \lambda_{f+1}$ . Let  $\rho \in {}^{\omega^\uparrow}\lambda_{f+1}$  with  $0\rho = \alpha$ . Since  $\Lambda_{f+1}^{[1,f+1]} \varphi \subseteq C_{f+1}$ , we can define a function  $\mathfrak{f}_\nu : \Lambda^{[1,f]} \rightarrow C_{f+1}$  for every  $\nu \in {}^{\omega^\uparrow}\lambda_{f+1}$  by  $\bar{\eta}' \mathfrak{f}_\nu = (\bar{\eta}' \wedge \nu) \varphi$ . This, in turn, gives us a function  $\mathfrak{f} : {}^{\omega^\uparrow}\lambda_{f+1} \rightarrow \mathfrak{P}$ . By Lemma 12, there exists  $\eta \in {}^{\omega^\uparrow}\lambda_{f+1}$  such that  $\rho \subset \eta$  and  $\Phi_{\eta n} = \mathfrak{f}_{\eta \restriction n}$  for all  $n < \omega$ .

Now we will use  $\varphi$  and  $\eta$  to define a function  $\varphi' : \Lambda_*^{[1,f]} \rightarrow C^f$ . For every  $\bar{\nu} \in \Lambda_*^{[1,f]}$ , define  $\bar{\nu} \varphi' = (\bar{\nu} \wedge \eta) \varphi$ ; and observe that  $\Lambda_m^{[1,f]} \varphi' \subseteq C_m$  for all  $m \in [1, f]$ . By induction hypothesis, there exists  $\bar{\eta}' \in \Lambda^{[1,f]}$  such that  $\varphi' \restriction [\bar{\eta}'] = \varphi_{\bar{\eta}'}^f$ . Define  $\bar{\eta} = \bar{\eta}' \wedge \eta$  to obtain  $\eta_{f+1} = \eta$ . Finally, we must verify that  $\varphi \restriction [\bar{\eta}] = \varphi_{\bar{\eta}}^{f+1}$ . If  $m \in [1, f]$ , then

$$(\bar{\eta} \restriction \langle m, n \rangle) \varphi = (\bar{\eta}' \restriction \langle m, n \rangle) \varphi' = (\bar{\eta}' \restriction \langle m, n \rangle) \varphi_{\bar{\eta}'}^f = (\bar{\eta} \restriction \langle m, n \rangle) \varphi_{\bar{\eta}}^{f+1},$$

and if  $m = f+1$ , then

$$(\bar{\eta} \restriction \langle m, n \rangle) \varphi = \bar{\eta}' \mathfrak{f}_{\eta_{f+1}n} = \bar{\eta}' \Phi_{\eta_{f+1}n} = (\bar{\eta} \restriction \langle m, n \rangle) \varphi_{\bar{\eta}}^{f+1}.$$

This completes the proof. □

## 2. The Construction

Let  $\bar{\lambda} = \langle \lambda_1, \dots, \lambda_k \rangle$  be a  $\blacksquare$ -sequence. We consider the sets  $\Lambda$  and  $\Lambda_*$  as before.

Given a subset  $Y_* \subseteq \Lambda_*$ , we consider the free abelian group

$$B_{Y_*} = \bigoplus_{\bar{\nu} \in Y_*} \mathbb{Z}e_{\bar{\nu}}.$$

If  $Y_* = [\bar{\eta}]$  for some  $\bar{\eta} \in \Lambda$ , then we simply write  $B_{\bar{\eta}}$ . The starting point of the final construction of this section will be the abelian group

$$B = \bigoplus_{\bar{\nu} \in \Lambda_*} \mathbb{Z}e_{\bar{\nu}}.$$

Moreover, let

$$\bar{B} = \hat{B} \cap \prod_{\bar{\nu} \in \Lambda_*} \mathbb{Z}e_{\bar{\nu}}$$

where  $\hat{B}$  denotes the  $p$ -completion of  $B$  for some prime  $p$ . Finally, let  $S = \bigoplus_{n < \omega} \mathbb{Z}e_n$ .

DEFINITION 16. For every  $b = \sum_{\bar{\nu} \in \Lambda_*} b_{\bar{\nu}}e_{\bar{\nu}} \in \hat{B}$  with  $b_{\bar{\nu}} \in J_p$ , the  $\Lambda_*$ -support of  $b$  is the set

$$[b] = \{ \bar{\nu} \mid b_{\bar{\nu}} \neq 0 \}.$$

DEFINITION 17. Let  $N$  be an abelian group. A *trap* for  $B$ ,  $N$  is a homomorphism

$$\varphi_{\bar{\eta}} : B_{\bar{\eta}} \rightarrow N.$$

THE  $\bar{\lambda}$ -BLACK BOX 18. Let  $\bar{\lambda} = \langle \lambda_1, \dots, \lambda_k \rangle$  be a  $\blacksquare$ -sequence,  $\Lambda$  and  $\Lambda_*$  as before and  $N$  an abelian group such that  $|N| \leq \chi_1$ . Then there exists a family of traps  $\langle \varphi_{\bar{\eta}} \mid \bar{\eta} \in \Lambda \rangle$  satisfying the following

Prediction Principle: If  $\varphi : B \rightarrow N$  is any homomorphism and  $\alpha < \lambda_k$ , then there exists  $\bar{\eta} \in \Lambda$  such that  $\varphi \upharpoonright B_{\bar{\eta}} = \varphi_{\bar{\eta}}$  and  $0\eta_k = \alpha$ .

PROOF. The members of the sequence  $\bar{C} = \langle C_1, \dots, C_k \rangle$  where  $C_m = N$  for all  $m \in [1, k]$  satisfy  $|C_m| \leq \chi_m$  since  $\bar{\lambda}$  is increasing and  $|N| \leq \chi_1$ . Hence,  $C = N$ . The  $\bar{\lambda}$ -Black Box 15 provides us with a family of set-traps  $\langle \varphi_{\bar{\eta}} \mid \bar{\eta} \in \Lambda \rangle$ . Since  $[\bar{\eta}] \subset \Lambda_*$ , each  $\varphi_{\bar{\eta}}$  can be regarded as a homomorphism  $\varphi_{\bar{\eta}} : B_{\bar{\eta}} \rightarrow N$ . Since any homomorphism  $\varphi : B \rightarrow N$  is completely determined by its action on the generators  $e_{\bar{\nu}}$  of  $B$ , it can be regarded as a function  $\varphi : \Lambda_* \rightarrow C$  which also satisfies  $\Lambda_m \varphi \subseteq C_m$ . Thus, for  $\alpha < \lambda_k$ , the  $\bar{\lambda}$ -Black Box 15 yields that there exists  $\bar{\eta} \in \Lambda$  such that  $0\eta_k = \alpha$  and  $\varphi \upharpoonright [\bar{\eta}] = \varphi_{\bar{\eta}}$ , i.e.  $\varphi \upharpoonright B_{\bar{\eta}} = \varphi_{\bar{\eta}}$ .  $\square$

DEFINITION 19. For  $\bar{\eta} \in \Lambda$  and  $n < \omega$ , we define the *branch element* associated with  $\bar{\eta}$  and  $n$  as

$$y_{\bar{\eta}n} = \sum_{i=n}^{\infty} p^{i-n} \left( \sum_{m=1}^k e_{\bar{\eta} \upharpoonright \langle m, i \rangle} \right).$$

We write  $y_{\bar{\eta}}$  for  $y_{\bar{\eta}0}$ . Choose an element  $b_{\bar{\eta}} = \sum_{i=0}^{\infty} p^i b_i^{\bar{\eta}} \in \bar{B}$ , where  $b_i^{\bar{\eta}} \in B$  for all  $i < \omega$ , and put  $b_{\bar{\eta}n} = \sum_{i=n}^{\infty} p^{i-n} b_i^{\bar{\eta}} \in \bar{B}$  (so  $b_{\bar{\eta}} = b_{\bar{\eta}0}$ ). We define the *branch-like element* associated with  $\bar{\eta}$  and  $n$  as

$$y'_{\bar{\eta}n} = b_{\bar{\eta}n} + y_{\bar{\eta}n}.$$

We also write  $y'_{\bar{\eta}}$  for  $y'_{\bar{\eta}0}$ .

Since  $|S| = \aleph_0 < \chi_1$ , the  $\bar{\lambda}$ -Black Box 18 ensures the existence of a family of traps  $\langle \varphi_{\bar{\eta}} \mid \bar{\eta} \in \Lambda \rangle$  such that, if  $\varphi : B \rightarrow S$  is any homomorphism and  $\alpha < \lambda_k$ , then there exists  $\bar{\eta} \in \Lambda$  with  $\varphi \upharpoonright B_{\bar{\eta}} = \varphi_{\bar{\eta}}$  and  $0\eta_k = \alpha$ .

For  $b \in \bar{B}$ , define  $\|b\| = \sup_{\bar{\nu} \in [b]} \|\bar{\nu}\|$ . For  $\alpha < \lambda_k$ , let  $\bar{B}_\alpha = \langle b \in \bar{B} \mid \|b\| < \alpha \rangle$ . We fix a map  $\delta : \lambda_k \rightarrow \bar{B}$  such that  $\alpha\delta \in \bar{B}_\alpha$  for all  $\alpha < \lambda_k$ .

**STEP LEMMA 20.** *Let  $\bar{\eta} \in \Lambda$ ,  $b_{\bar{\eta}} = 0\eta_k\delta$  and  $\varphi_{\bar{\eta}}$  from the  $\bar{\lambda}$ -Black Box 18. There exists an  $\varepsilon_{\bar{\eta}} \in \{0, 1\}$  such that no homomorphism  $\varphi : \langle B, y'_{\bar{\eta}} = \varepsilon_{\bar{\eta}}b_{\bar{\eta}} + y_{\bar{\eta}} \rangle_* \rightarrow S$  satisfies both  $\varphi \upharpoonright B_{\bar{\eta}} = \varphi_{\bar{\eta}}$  and  $b_{\bar{\eta}}\varphi \in \widehat{S} \setminus S$ .*

**PROOF.** Suppose towards a contradiction that for both  $\varepsilon \in \{0, 1\}$ , there exists some  $\varphi^\varepsilon : \langle B, \varepsilon b_{\bar{\eta}} + y_{\bar{\eta}} \rangle_* \rightarrow S$  such that  $\varphi^\varepsilon \upharpoonright B_{\bar{\eta}} = \varphi_{\bar{\eta}}$  and  $b_{\bar{\eta}}\varphi^\varepsilon \in \widehat{S} \setminus S$ . On one hand,  $(b_{\bar{\eta}} + y_{\bar{\eta}})\varphi^1 - y_{\bar{\eta}}\varphi^0 \in S$ . On the other hand,  $(b_{\bar{\eta}} + y_{\bar{\eta}})\varphi^1 - y_{\bar{\eta}}\varphi^0 = b_{\bar{\eta}}\varphi^1 \in \widehat{S} \setminus S$ , which is the desired contradiction.  $\square$

Let  $\mathfrak{F} = \{ y'_{\bar{\eta}} = \varepsilon_{\bar{\eta}}b_{\bar{\eta}} + y_{\bar{\eta}} \mid \bar{\eta} \in \Lambda, b_{\bar{\eta}} = 0\eta_k\delta \}$  be the family of branch-like elements obtained after choosing every  $\varepsilon_{\bar{\eta}}$  by means of the Step Lemma 20. Define

$$G = \langle B, y'_{\bar{\eta}n} \mid y'_{\bar{\eta}} \in \mathfrak{F}, n < \omega \rangle = \langle B, y'_{\bar{\eta}} \mid y'_{\bar{\eta}} \in \mathfrak{F} \rangle_*.$$

Notice the following divisibility property of the branch-like elements:

$$py'_{\bar{\eta}(n+1)} = y'_{\bar{\eta}n} - \sum_{m=1}^k e_{\bar{\eta} \setminus (m,n)} - \varepsilon_{\bar{\eta}} b_{\bar{\eta}} \in G.$$

Since  $G \subseteq_* \bar{B} \subseteq \prod_{\bar{\nu} \in \Lambda_*} \mathbb{Z}e_{\bar{\nu}}$ , the group  $G$  is separable.

DEFINITION 21. For  $g \in G$ , define the  $\Lambda$ -support  $[g]_\Lambda$  of  $g$  to be the set of elements of  $\Lambda$  that contribute to the representation of  $g$ . More precisely, if  $p^m g = b + \sum_{\bar{\eta} \in \Lambda} n_{\bar{\eta}} y'_{\bar{\eta}}$  for some  $m \geq 0$ , where  $b \in B$  and  $n_{\bar{\eta}} \in \mathbb{Z}$  for all  $\bar{\eta} \in \Lambda$ , then

$$[g]_\Lambda = \{ \bar{\eta} \in \Lambda \mid n_{\bar{\eta}} \neq 0 \}.$$

Obviously,  $[g]_\Lambda$  is finite. For  $H \subseteq G$ , we define  $[H]_\Lambda = \bigcup_{g \in H} [g]_\Lambda$ .

DEFINITION 22. A module  $N$  is  $\kappa$ -free if every subset of  $N$  of cardinality  $< \kappa$  is contained in a free submodule of  $N$ .

THEOREM 23. The group  $G$  defined as above is  $\aleph_k$ -free.

PROOF. Suppose that  $H$  is a subset of  $G$  of cardinality  $\aleph_{k-1}$ . Let  $\sigma : \bar{B} \rightarrow [\Lambda_*]^{\leq \aleph_0}$  be the “ $\Lambda_*$ -support” function, i.e.  $b\sigma = [b]$  for all  $b \in \bar{B}$ . Notice that  $F : \Lambda \rightarrow [\Lambda_*]^{\leq \aleph_0}$  given by  $\bar{\eta}F = 0\eta_k \delta \sigma$  is regressive since  $\alpha \delta \in \bar{B}_\alpha$  for all  $\alpha < \lambda_k$ . Let  $\Omega = [H]_\Lambda$ ,  $\Omega_* = [H] \setminus ([\Omega] \cup \bigcup_{\bar{\eta} \in \Omega} [\bar{\eta}F])$  and observe that  $|\Omega| \leq \aleph_{k-1}$ . Then the subgroups

$$G_\Omega = \langle e_{\bar{\eta} \langle m, n \rangle}, e_{\bar{\nu}}, y'_{\bar{\eta}} \mid \bar{\eta} \in \Omega, \bar{\nu} \in \bar{\eta}F, m \in [1, k], n < \omega \rangle_*,$$

and

$$G' = G_\Omega \oplus B_{\Omega_*}$$

satisfy  $H \subseteq G'$ . Our goal is to show that  $G'$  is free, for which it suffices to show that  $G_\Omega$  is free. Suppose that  $|\Omega| = \aleph_{k-1}$ . By taking  $u_{\bar{\eta}} = [1, k]$  for all  $\bar{\eta} \in \Omega$ , we enumerate

$\Omega = \{ \bar{\eta}^\alpha \mid \alpha < \zeta \}$  for some  $\zeta \in [\omega_{k-1}, \omega_k)$  according to the Freeness Proposition 11 and find  $\ell_\alpha \in u_{\bar{\eta}}$  and  $n_\alpha < \omega$  such that

$$\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle \notin \{ \bar{\eta}^\beta \restriction \langle \ell_\alpha, n \rangle \mid \beta < \alpha \} \cup \bigcup \Omega_\alpha F$$

for all  $n \geq n_\alpha$ . This allows us to write

$$G_\Omega = \langle e_{\bar{\eta}^\alpha \restriction \langle m, n \rangle}, e_{\bar{\nu}}, y'_{\bar{\eta}^\alpha n} \mid \alpha < \zeta, \bar{\nu} \in \bar{\eta}^\alpha F, m \in [1, k], n < \omega \rangle.$$

Let

$$G_\alpha = \langle e_{\bar{\eta}^\beta \restriction \langle m, n \rangle}, e_{\bar{\nu}}, y'_{\bar{\eta}^\beta n} \mid \beta < \alpha, \bar{\nu} \in \bar{\eta}^\beta F, m \in [1, k], n < \omega \rangle$$

and notice that  $G_0 = 0$ ,  $\bigcup_{\alpha < \zeta} G_\alpha = G_\Omega$  and

$$\begin{aligned} G_{\alpha+1} &= G_\alpha + \langle e_{\bar{\eta}^\alpha \restriction \langle m, n \rangle}, e_{\bar{\nu}}, y'_{\bar{\eta}^\alpha n} \mid \bar{\nu} \in \bar{\eta}^\alpha F, m \in [1, k], n < \omega \rangle \\ &= G_\alpha + \langle e_{\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle} \mid n < n_\alpha \rangle + \langle y'_{\bar{\eta}^\alpha n} \mid n \geq n_\alpha \rangle + \langle e_{\bar{\nu}} \mid \bar{\nu} \in \bar{\eta}^\alpha F \rangle \\ &\quad + \langle e_{\bar{\eta}^\alpha \restriction \langle m, n \rangle} \mid m \in [1, k] \setminus \{\ell_\alpha\}, n < \omega \rangle \end{aligned}$$

since

$$e_{\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle} = y'_{\bar{\eta}^\alpha n} - p y'_{\bar{\eta}^\alpha (n+1)} - b_n^{\bar{\eta}^\alpha} - \sum_{\substack{m=1 \\ m \neq \ell_\alpha}}^k e_{\bar{\eta}^\alpha \restriction \langle m, n \rangle}$$

for  $n \geq n_\alpha$  and

$$y'_{\bar{\eta}^\alpha n} = p^{n_\alpha - n} y'_{\bar{\eta}^\alpha n_\alpha} + \sum_{i=0}^{n_\alpha - n - 1} p^i \left( b_{n+i}^{\bar{\eta}^\alpha} + \sum_{m=1}^k e_{\bar{\eta}^\alpha \restriction \langle m, n+i \rangle} \right)$$

for  $n < n_\alpha$ . We claim that  $G_{\alpha+1}/G_\alpha$  is free. To prove our claim, suppose

$$\underbrace{\sum_{n < n_\alpha} z_n e_{\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle}}_{(1)} + \underbrace{\sum_{n \geq n_\alpha} z_n y'_{\bar{\eta}^\alpha n}}_{(2)} + \underbrace{\sum_{n < \omega} \sum_{\substack{m=1 \\ m \neq \ell_\alpha}}^k z_{mn} e_{\bar{\eta}^\alpha \restriction \langle m, n \rangle}}_{(3)} + \underbrace{\sum_{\bar{\nu} \in \bar{\eta}^\alpha F} z_{\bar{\nu}} e_{\bar{\nu}}}_{(4)} \in G_\alpha$$



It is immediate that the support of term (1) is disjoint from those of the other terms. Then  $z_n = 0$  for all  $n < n_\alpha$  with  $e_{\bar{\eta}^\alpha \upharpoonright \langle \ell_\alpha, n \rangle} \notin G_\alpha$ . By the Freeness Proposition 11,  $\bar{\eta}^\alpha \upharpoonright \langle \ell_\alpha, n \rangle$  neither belongs to the support of the terms (3) and (4) nor to the support of  $G_\alpha$  for all  $n \geq n_\alpha$ , which implies that  $z_n = 0$  for all  $n \geq n_\alpha$ . Since  $\|b_{\bar{\eta}^\alpha}\| < 0\eta_k^\alpha$ , it follows that  $z_{mn} = z_{\bar{\nu}} = 0$  for all  $m \in [1, k] \setminus \{\ell_\alpha\}$ ,  $n < \omega$  and  $\bar{\nu} \in \bar{\eta}^\alpha F$  such that  $e_{\bar{\eta}^\alpha \upharpoonright \langle m, n \rangle}, e_{\bar{\nu}} \notin G_\alpha$ . Therefore,  $G_{\alpha+1}/G_\alpha$  is freely generated by the set

$$\{e_{\bar{\eta}^\alpha \upharpoonright \langle \ell_\alpha, r \rangle}, y'_{\bar{\eta}^\alpha s}, e_{\bar{\eta}^\alpha \upharpoonright \langle m, t \rangle}, e_{\bar{\nu}} \mid$$

$$r < n_\alpha, s \geq n_\alpha, m \in [1, k] \setminus \{\ell_\alpha\}, t < \omega, \bar{\nu} \in \bar{\eta}^\alpha F\} \setminus G_\alpha.$$

Since  $G_\Omega$  is the union of the continuous chain  $\{G_\alpha \mid \alpha < \zeta\}$  such that  $G_0$  is free and every  $G_{\alpha+1}/G_\alpha$  is free,  $G_\Omega$  itself is free. We conclude that  $G$  is  $\aleph_k$ -free.  $\square$

**THEOREM 24.** *For any  $\blacksquare$ -sequence  $\langle \lambda_1, \dots, \lambda_k \rangle$  such that  $\chi_k = \lambda_k$ , there exists a separable  $\aleph_k$ -free abelian group  $G$  with no epimorphisms onto  $S$ .*

**PROOF.** Let  $G$  be as before. Notice that the additional condition  $\chi_k = \lambda_k$  implies that we can fix the map  $\delta : \lambda_k \rightarrow \bar{B}$  to be surjective. Suppose  $\varphi : G \rightarrow S$  is an epimorphism. If  $\bar{B}\varphi \subseteq S$ , then  $e_{\bar{\nu}}\varphi \neq 0$  for only finitely many  $\bar{\nu} \in \Lambda_*$ . Hence,  $\varphi$  cannot be an epimorphism. It follows that  $\bar{B}\varphi \cap \hat{S} \setminus S \neq \emptyset$ . Take any  $g \in \bar{B}$  such that  $g\varphi = s \in \hat{S} \setminus S$ . Then there exists  $\alpha < \lambda_k$  such that  $\alpha\delta = g$ . By the  $\bar{\lambda}$ -Black Box 18, there exists  $\bar{\eta} \in \Lambda$  such that  $\varphi \upharpoonright B_{\bar{\eta}} = \varphi_{\bar{\eta}}$  and  $0\eta_k = \alpha$ . But then  $\psi = \varphi \upharpoonright \langle B, y'_{\bar{\eta}} \rangle_*$  satisfies both  $\psi \upharpoonright B_{\bar{\eta}} = \varphi_{\bar{\eta}}$  and  $b_{\bar{\eta}}\psi = b_{\bar{\eta}}\varphi = (0\eta_k\delta)\varphi = (\alpha\delta)\varphi = g\varphi = s \in \hat{S} \setminus S$ , contradicting the choice of  $\varepsilon_{\bar{\eta}}$ . Therefore  $G$  has no epimorphisms onto  $S$ .  $\square$

This theorem implies that every time we have a homomorphism  $\varphi : G \rightarrow H$  with  $H$  a free abelian group, then  $G\varphi \subseteq H$  will have finite rank.

### 3. Slenderness

We have constructed a separable  $\aleph_k$ -free group  $G$  satisfying the property stated at the end of the previous section. We now show that this group is slender. Let  $P = \prod_{n < \omega} \mathbb{Z}e_n$ .

DEFINITION 25. A torsion-free group  $G$  is *slender* if for all homomorphism  $\varphi : P \rightarrow G$ , we have  $e_n\varphi = 0$  for almost all  $n \in \omega$ .

The next theorem completely characterizes slender groups as those not containing certain groups as subgroups. See Fuchs [10] vol. 2, Theorem 95.3 for a proof.

THEOREM 26 (Nunke). *A torsion-free group is slender if and only if it contains no copy of  $\mathbb{Q}$ ,  $P$  or  $J_p$  for any prime  $p$ .*

Assume for the moment that the next Lemma is true.

LEMMA 27. *There exists a non-free subgroup of  $P$  of cardinality  $\aleph_1$ .*

THEOREM 28. *For  $k > 1$ , every  $\aleph_k$ -free group is slender.*

PROOF. Let  $G$  be an  $\aleph_k$ -free group for some  $k > 1$ . Since  $\mathbb{Q}$  is not free and  $|\mathbb{Q}| = \aleph_0$ ,  $G$  cannot contain copies of  $\mathbb{Q}$ . Since  $\mathbb{Z}(p) \subseteq J_p$  is not free,  $G$  cannot contain copies of  $J_p$ . By Lemma 27,  $P$  has a non-free subgroup of cardinality  $\aleph_1$ , so  $G$  does not contain a copy of  $P$  either. Thus by Theorem 26,  $G$  is slender.  $\square$

It only remains to prove Lemma 27. First we introduce some definitions. Let  $R$  be a commutative ring with 1 and  $\mathbb{S} \subseteq R \setminus \{0\}$  be multiplicatively closed.

DEFINITION 29. An  $R$ -module  $M$  is

- (1)  $\mathbb{S}$ -torsion-free if  $sm = 0$  implies  $m = 0$  for all  $s \in \mathbb{S}$ ,  $m \in M$ ;
- (2)  $\mathbb{S}$ -reduced if  $\bigcap_{s \in \mathbb{S}} sM = \{0\}$ ;
- (3)  $\mathbb{S}$ -divisible if  $sM = M$  for all  $s \in \mathbb{S}$ .
- (4) A submodule  $N$  of an  $R$ -module  $M$  is  $\mathbb{S}$ -dense if  $(m + sM) \cap N \neq \emptyset$  for all  $m \in M$ ,  $s \in \mathbb{S}$ .

LEMMA 30. A submodule  $N$  of an  $R$ -module  $M$  is  $\mathbb{S}$ -dense if and only if  $M/N$  is  $\mathbb{S}$ -divisible.

PROOF. For  $m \in M$ ,  $s \in \mathbb{S}$ , we have  $(m + sM) \cap N \neq \emptyset$  if and only if there exist  $m' \in M$ ,  $n \in N$  such that  $m - sm' = n$ . But, this holds exactly if  $m + N = sm' + N$ , i.e.  $M/N \subseteq s(M/N)$ .  $\square$

Now let  $R = \mathbb{Z}$ ,  $\mathbb{S} = \mathbb{Z}^+$  (the set of positive integers) and  $S$  as in the previous section.

DEFINITION 31. The subgroup

$$\overline{S} = \{ y \in P \mid \forall n \in \mathbb{Z}^+ ((y + nP) \cap S \neq \emptyset) \}$$

of  $P$  is called the  $\mathbb{S}$ -closure of  $S$ , which is the largest subgroup of  $P$  containing  $S$  as an  $\mathbb{S}$ -dense subgroup.

LEMMA 32. Let  $S \leq G \leq P$ . Then  $G = \overline{S}$  if and only if  $P/G$  is  $\mathbb{S}$ -reduced and  $G/S$  is  $\mathbb{S}$ -divisible.

PROOF. ( $\Rightarrow$ ) Let  $x \in P$  such that  $(x + \bar{S}) \in \bigcap_{n \in \mathbb{Z}^+} n(P/\bar{S})$ . Then for all  $n \in \mathbb{Z}^+$ , there exists  $y_n \in P$  such that  $x + \bar{S} = ny_n + \bar{S}$ , so  $x - ny_n \in \bar{S}$ . Then for all  $m, n \in \mathbb{Z}^+$ ,  $((x - ny_n) + mP) \cap S \neq \emptyset$ . In particular,  $((x - ny_n) + nP) \cap S = (x + nP) \cap S \neq \emptyset$ , which implies that  $x \in \bar{S}$ . Hence,  $P/G$  is  $\mathbb{S}$ -reduced. Since  $S$  is a  $\mathbb{S}$ -dense subgroup of  $\bar{S}$ , it follows that  $\bar{S}/S$  is  $\mathbb{S}$ -divisible by Lemma 30.

( $\Leftarrow$ ) By Lemma 30,  $S$  is  $\mathbb{S}$ -dense in  $G$ , so  $G \leq \bar{S}$ . Let  $x \in \bar{S}$ . For all  $n \in \mathbb{Z}^+$ ,  $(x + nP) \cap S \neq \emptyset$ . Thus, there exist  $y_n \in P$  such that  $x - ny_n \in S \subset G$ . It follows that  $x + G = ny_n + G \in \bigcap_{n \in \mathbb{Z}^+} n(P/G)$ , which implies that  $x \in G$  because  $P/G$  is  $\mathbb{S}$ -reduced. Therefore,  $\bar{S} \leq G$ .  $\square$

The next lemma follows immediately from Fuchs [10], Vol. 1, Theorem 23.1 on the structure of divisible groups.

LEMMA 33. *If  $G$  is torsion-free and divisible, then  $G$  is a  $\mathbb{Q}$ -vector space and  $G \cong \bigoplus_{i \in I} \mathbb{Q}e_i$  for some  $I$  such that  $|I| = \dim_{\mathbb{Q}} G$ .*

THEOREM 34.  *$\bar{S}/S$  is a  $\mathbb{Q}$ -vector space and  $\text{rk } \bar{S}/S = \dim_{\mathbb{Q}} \bar{S}/S = 2^{\aleph_0}$ .*

PROOF.  $\bar{S}/S$  is torsion-free and divisible, so by Lemma 33,  $\bar{S}/S$  is a  $\mathbb{Q}$ -vector space. Moreover,  $|S| = \aleph_0$  and  $|\bar{S}| = 2^{\aleph_0}$ , so  $2^{\aleph_0} = |\bar{S}| = |S| \cdot |\bar{S}/S| = \aleph_0 \cdot |\bar{S}/S|$ . Therefore,  $\text{rk } \bar{S}/S = \dim_{\mathbb{Q}} \bar{S}/S = 2^{\aleph_0}$ .  $\square$

PROPOSITION 35. *Let  $F$  be a free abelian group and let  $H \neq 0$  be a subgroup of  $F$  such that  $|H| < |F|$ . Then there exist (free)  $C_1, C_2$  such that  $F = C_1 \oplus C_2$ ,  $H \subseteq C_1$  and  $|C_2| = |F|$ .*

PROOF. Let  $B = \{e_i \mid i \in I\}$  be a basis of  $F$ . Every element  $f \in F$  can be written as an infinite sum  $\sum_{i \in I} n_i e_i$  where  $n_i \in \mathbb{Z}$  and  $n_i = 0$  for almost all  $i \in I$ . For every  $f \in F$ , we consider the support  $[f] = \{e_i \mid n_i \neq 0\}$  of  $f$ , which is a finite subset of  $B$ . Put  $B_1 = \bigcup_{f \in H} [f]$  and  $B_2 = B \setminus B_1$ . Define  $C_i = \langle B_i \rangle$  for  $i = 1, 2$ . By construction,  $C_1 \cap C_2 = \{0\}$  and  $H \subseteq C_1$ . Since  $|B_1| = |H|$ , we conclude  $|B_2| = |C_2| = |F|$ .  $\square$

*Proof of Lemma 27.* By Theorem 34,  $\overline{S}/S$  is a  $\mathbb{Q}$ -vector space with  $\dim_{\mathbb{Q}} \overline{S}/S = 2^{\aleph_0}$ . Thus there exists a subspace  $V' \leq \overline{S}/S$  such that  $\dim_{\mathbb{Q}} V' = \aleph_1$ . There exists some  $S \leq V \leq \overline{S} \leq P$  such that  $V/S \cong V'$ . Since  $|S| = \aleph_0$ , we have  $|V| = \aleph_1$ . We claim that  $V$  is not free. Suppose towards a contradiction that  $V$  is free. By Proposition 35, there exist some free  $C_1, C_2$  such that  $V = C_1 \oplus C_2$ ,  $S \subseteq C_1$  and  $|C_2| = |V|$ . By construction  $S \cap C_2 = \{0\}$ , so that  $V' \cong C_1/S \oplus C_2$  is a  $\mathbb{Q}$ -vector space with a free direct summand  $C_2$ , a contradiction. Therefore  $V$  is not free.  $\square$

## CHAPTER 2

### Separable $\aleph_k$ -free Groups Satisfying $\text{End } G = A \oplus \text{Fin } G$

#### 1. Set-theoretic Modifications

Let  $A$  be a ring with free additive group  $A^+ = \bigoplus_{\alpha < \kappa} \mathbb{Z}e_\alpha$  such that

$$\overline{A} = \widehat{A} \cap \prod_{\alpha < \kappa} \mathbb{Z}e_\alpha$$

is an  $A$ -module, where  $\widehat{A}$  denotes the  $p$ -completion of  $A$  for some prime  $p$ . Such rings are called *separably realizable*. It was shown at different times and independently that  $\overline{A}$  is a ring for  $\kappa = \aleph_0$  by Goodearl, Menal, Moncasi [15], Corner, Göbel [3] and Nielsen [19]. For more details, see Göbel, Trlifaj [14].

We recursively construct a sequence of infinite cardinals  $\langle \lambda_1, \dots, \lambda_k \rangle$  as follows:

- (1) Let  $\lambda_0 = |A|$ .
- (2) Suppose we have constructed  $\lambda_m$  for some  $m \in [0, k)$ . Choose some cardinal

$$\mu_{m+1} \text{ such that } \mu_{m+1}^{\lambda_m} = \mu_{m+1} \text{ and let } \lambda_{m+1} = \mu_{m+1}^+.$$

Once we have the cardinal sequence  $\langle \lambda_1, \dots, \lambda_k \rangle$ , we construct the sets  $\Lambda$  and  $\Lambda_*$  as before. Given a subset  $Y_* \subseteq \Lambda_*$ , we consider the free  $A$ -module

$$B_{Y_*} = \bigoplus_{\overline{\nu} \in Y_*} Ae_{\overline{\nu}}.$$

The basic  $A$ -module on which we base the final construction is the free  $A$ -module

$$B = \bigoplus_{\bar{\nu} \in \Lambda_*} A e_{\bar{\nu}}.$$

Moreover, let

$$\bar{B} = \widehat{B} \cap \prod_{\bar{\nu} \in \Lambda_*} \bar{A} e_{\bar{\nu}}$$

where  $\widehat{B}$  denotes the  $p$ -completion of  $B$ .

Due to the existence of non-hereditary rings, it is necessary to modify the notion of  $\kappa$ -free modules. The following definition of  $\kappa$ -freeness is due to Göbel, Herden, Shelah [11], which is a slightly stronger version of the one in Eklof, Mekler [7].

**DEFINITION 36.** If  $\kappa$  is a regular uncountable cardinal, we say that an  $A$ -module  $M$  is  $\kappa$ -free if there is a family  $\mathcal{C}$  of  $p$ -pure  $A$ -submodules of  $M$  satisfying:

- (1) every element of  $\mathcal{C}$  is  $< \kappa$ -generated and free;
- (2) every element of  $[M]^{<\kappa}$  is contained in an element of  $\mathcal{C}$ ;
- (3)  $\mathcal{C}$  is closed under unions of well-ordered chains of length  $< \kappa$ .

**DEFINITION 37.** Let  $Y_* \subseteq \Lambda_*$ ,  $Y \subseteq \Lambda$  and  $\mathfrak{F}_{Y_*Y} = \{ y'_{\bar{\eta}} = b_{\bar{\eta}} + y_{\bar{\eta}} \mid \bar{\eta} \in Y, b_{\bar{\eta}} \in \bar{B}_{Y_*} \}$  be a family of branch-like elements.

- (1) If  $X \subseteq Y$ , then  $\mathfrak{F}_{Y_*X} = \{ y'_{\bar{\eta}} \in \mathfrak{F}_{Y_*Y} \mid \bar{\eta} \in X \}$ .
- (2) We say that  $\mathfrak{F}_{Y_*Y}$  is *regressive* if  $\| b_{\bar{\eta}} \| < 0\eta_k$  for all  $\bar{\eta} \in Y$ .
- (3) We say that the pair  $(Y_*, Y)$  is  $\Lambda$ -closed if for all  $\bar{\eta} \in Y$  there exists some (minimal)  $N_{\bar{\eta}} < \omega$  such that  $[\bar{\eta}]_{N_{\bar{\eta}}} \subseteq Y_*$  (see Definition 7).



- (4) If  $(Y_*, Y)$  is  $\Lambda$ -closed and  $\mathfrak{F}_{Y_*Y}$  is a regressive family of branch-like elements, we define the  $A$ -module

$$G_{Y_*Y} = \langle B_{Y_*}, Ay'_{\bar{\eta}n} \mid \bar{\eta} \in Y, n \geq N_{\bar{\eta}} \rangle = \langle B_{Y_*}, Ay'_{\bar{\eta}N_{\bar{\eta}}} \mid \bar{\eta} \in Y \rangle_*.$$

OBSERVATION 38. If  $G_{Y_*Y}$  is defined as before,  $X \subseteq Y$ ,  $X_* \subseteq Y_*$ ,  $(X_*, X)$  is a  $\Lambda$ -closed pair, and if  $b_{\bar{\eta}} \in \overline{B}_{X_*}$  for all  $\bar{\eta} \in X$ , then the  $A$ -module  $G_{X_*X}$  generated by  $\mathfrak{F}_{X_*X} = \mathfrak{F}_{Y_*X}$  is an  $A$ -submodule of  $G_{Y_*Y}$ .

THEOREM 39. *If  $(Y_*, Y)$  is  $\Lambda$ -closed and  $\mathfrak{F}_{Y_*Y}$  is a regressive family of branch-like elements, then  $G_{Y_*Y}$  is an  $\aleph_k$ -free  $A$ -module.*

PROOF. We will construct a family  $\mathcal{C}$  of  $p$ -pure  $A$ -submodules of  $G_{Y_*Y}$  according to Definition 36. For every  $H \in [G_{Y_*Y}]^{<\aleph_k}$ , let  $\Omega = [H]_\Lambda$  and

$$\Omega_* = \left( [H] \cup \bigcup_{\bar{\eta} \in \Omega} [\bar{\eta}] \cup [b_{\bar{\eta}}] \right) \cap Y_*.$$

Then  $[\bar{\eta}]_{N_{\bar{\eta}}} \subseteq \Omega_*$  for all  $\bar{\eta} \in \Omega$ . It follows that  $(\Omega_*, \Omega)$  is  $\Lambda$ -closed, so together with  $\mathfrak{F}_{\Omega_*\Omega}$ , we can define the module  $G_{\Omega_*\Omega}$  which contains  $H$ . Observe that  $G_{\Omega_*\Omega}$  is  $<\aleph_k$ -generated because  $|\Omega|, |\Omega_*| \leq |H| \cdot \aleph_0 < \aleph_k$ . Our goal is to prove that  $G_{\Omega_*\Omega}$  is free. For that purpose, let  $\Omega'_* = \bigcup_{\bar{\eta} \in \Omega} [\bar{\eta}]_{N_{\bar{\eta}}} \cup [b_{\bar{\eta}N_{\bar{\eta}}}] \subseteq \Omega_*$ . Since  $Ae_{\bar{\nu}}$  is a free direct summand of  $G_{\Omega_*\Omega}$  for every  $\bar{\nu} \in \Omega_* \setminus \Omega'_*$ , it is enough to show that  $G_{\Omega'_*\Omega}$  is free.

Suppose that  $|\Omega| = \aleph_{k-1}$ . Let  $F : \Lambda \rightarrow [\Lambda_*]^{\leq \aleph_0}$  be any regressive map such that  $\bar{\eta}F = [b_{\bar{\eta}}]$  for all  $\bar{\eta} \in \Omega$ . By taking  $u_{\bar{\eta}} = [1, k]$  for all  $\bar{\eta} \in \Omega$ , we enumerate  $\Omega = \{\bar{\eta}^\alpha \mid \alpha < \zeta\}$  for some  $\zeta \in [\omega_{k-1}, \omega_k)$  according to the Freeness Proposition 11

and find  $\ell_\alpha \in u_{\bar{\eta}}$  and  $n_\alpha < \omega$  such that  $\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle \notin \{ \bar{\eta}^\beta \restriction \langle \ell_\alpha, n \rangle \mid \beta < \alpha \} \cup \bigcup \Omega_\alpha F$  for all  $n \geq n_\alpha$ . This allows us to write

$$G_{\Omega'_* \Omega} = \langle Ae_{\bar{\eta}^\alpha \restriction \langle m, n \rangle}, Ae_{\bar{\nu}}, Ay'_{\bar{\eta}^\alpha n} \mid \alpha < \zeta, \bar{\nu} \in [b_{\bar{\eta}^\alpha N_{\bar{\eta}^\alpha}}] \subseteq \bar{\eta}^\alpha F, m \in [1, k], n \in [N_{\bar{\eta}^\alpha}, \omega) \rangle.$$

For convenience, we also take  $n_\alpha > N_{\bar{\eta}^\alpha}$ . Let

$$G_\alpha = \langle Ae_{\bar{\eta}^\beta \restriction \langle m, n \rangle}, Ae_{\bar{\nu}}, Ay'_{\bar{\eta}^\beta n} \mid \beta < \alpha, \bar{\nu} \in [b_{\bar{\eta}^\beta N_{\bar{\eta}^\beta}}] \subseteq \bar{\eta}^\beta F, m \in [1, k], n \in [N_{\bar{\eta}^\beta}, \omega) \rangle$$

and notice that  $G_0 = 0$ ,  $\bigcup_{\alpha < \zeta} G_\alpha = G_{\Omega'_* \Omega}$  and

$$\begin{aligned} G_{\alpha+1} &= G_\alpha + \langle Ae_{\bar{\eta}^\alpha \restriction \langle m, n \rangle}, Ae_{\bar{\nu}}, Ay'_{\bar{\eta}^\alpha n} \mid \bar{\nu} \in [b_{\bar{\eta}^\alpha N_{\bar{\eta}^\alpha}}] \subseteq \bar{\eta}^\alpha F, m \in [1, k], n \in [N_{\bar{\eta}^\alpha}, \omega) \rangle \\ &= G_\alpha + \langle Ae_{\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle} \mid n \in [N_{\bar{\eta}^\alpha}, n_\alpha) \rangle + \langle Ay'_{\bar{\eta}^\alpha n} \mid n \geq n_\alpha \rangle \\ &\quad + \langle Ae_{\bar{\eta}^\alpha \restriction \langle m, n \rangle} \mid m \in [1, k] \setminus \{\ell_\alpha\}, n \in [N_{\bar{\eta}^\alpha}, \omega) \rangle + \langle Ae_{\bar{\nu}} \mid \bar{\nu} \in [b_{\bar{\eta}^\alpha N_{\bar{\eta}^\alpha}}] \subseteq \bar{\eta}^\alpha F \rangle \end{aligned}$$

since

$$e_{\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle} = y'_{\bar{\eta}^\alpha n} - p y'_{\bar{\eta}^\alpha (n+1)} - b_n^{\bar{\eta}^\alpha} - \sum_{\substack{m=1 \\ m \neq \ell_\alpha}}^k e_{\bar{\eta}^\alpha \restriction \langle m, n \rangle}$$

for  $n \geq n_\alpha$  and

$$y'_{\bar{\eta}^\alpha n} = p^{n_\alpha - n} y'_{\bar{\eta}^\alpha n_\alpha} + \sum_{i=0}^{n_\alpha - n - 1} p^i \left( b_{n+i}^{\bar{\eta}^\alpha} + \sum_{m=1}^k e_{\bar{\eta}^\alpha \restriction \langle m, n+i \rangle} \right)$$

for  $n \in [N_{\bar{\eta}^\alpha}, n_\alpha)$ . We claim that  $G_{\alpha+1} = G_\alpha \oplus \bigoplus_{e \in G'_\alpha} Ae$  for some  $G'_\alpha$ . To prove our claim, suppose

$$\underbrace{\sum_{n \in [N_{\bar{\eta}^\alpha}, n_\alpha)} a_n e_{\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle}}_{(1)} + \underbrace{\sum_{n \geq n_\alpha} a_n y'_{\bar{\eta}^\alpha n}}_{(2)} + \underbrace{\sum_{n \in [N_{\bar{\eta}^\alpha}, \omega)} \sum_{\substack{m=1 \\ m \neq \ell_\alpha}}^k a_{mn} e_{\bar{\eta}^\alpha \restriction \langle m, n \rangle}}_{(3)} + \underbrace{\sum_{\bar{\nu} \in [b_{\bar{\eta}^\alpha N_{\bar{\eta}^\alpha}}]} a_{\bar{\nu}} e_{\bar{\nu}}}_{(4)} \in G_\alpha$$

It is immediate that the support of the term (1) is disjoint from those of the other terms.

Then  $a_n = 0$  for all  $n \in [N_{\bar{\eta}^\alpha}, n_\alpha)$  with  $e_{\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle} \notin G_\alpha$ . By the Freeness Proposition 11,

$\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle$  neither belongs to the support of the terms (3) and (4) nor to the support of  $G_\alpha$  for all  $n \geq n_\alpha$ , which implies that  $a_n = 0$  for all  $n \geq n_\alpha$ . Since  $\|b_{\bar{\eta}^\alpha}\| < 0\eta_k^\alpha$ , it follows that  $a_{mn} = a_{\bar{\nu}} = 0$  for all  $m \in [1, k] \setminus \{\ell_\alpha\}$ ,  $n \in [N_{\bar{\eta}^\alpha}, \omega)$  and  $\bar{\nu} \in [b_{\bar{\eta}^\alpha} N_{\bar{\eta}^\alpha}]$  such that  $e_{\bar{\eta}^\alpha \restriction \langle m, n \rangle}, e_{\bar{\nu}} \notin G_\alpha$ . Therefore  $G_{\alpha+1} = G_\alpha \oplus \bigoplus_{e \in G'_\alpha} Ae$  is free with

$$G'_\alpha = \{ e_{\bar{\eta}^\alpha \restriction \langle \ell_\alpha, r \rangle}, y'_{\bar{\eta}^\alpha s}, e_{\bar{\eta}^\alpha \restriction \langle m, t \rangle}, e_{\bar{\nu}} \mid$$

$$r \in [N_{\bar{\eta}^\alpha}, n_\alpha), s \geq n_\alpha, m \in [1, k] \setminus \{\ell_\alpha\}, t \in [N_{\bar{\eta}^\alpha}, \omega), \bar{\nu} \in [b_{\bar{\eta}^\alpha} N_{\bar{\eta}^\alpha}] \} \setminus G_\alpha,$$

which in turn implies that  $G_{\Omega'_* \Omega}$  is free. By setting  $\mathcal{C} = \{ G_{\Omega_* \Omega} \mid |\Omega|, |\Omega_*| < \aleph_k \}$ , we conclude that  $G_{Y_* Y}$  is  $\aleph_k$ -free.  $\square$

## 2. Triple Modules

In this section we introduce modules defined by triples of subsets of  $\Lambda$  and  $\Lambda_*$ . These triple modules have important freeness properties.

DEFINITION 40. Let  $X_* \subseteq Y_* \subseteq \Lambda_*$ ,  $Y \subseteq \Lambda$  and  $f \in [1, k]$ .

- (1) Define  $Y_{X_*} = \{\bar{\eta} \in Y \mid \exists n < \omega ([\bar{\eta}]_n \subseteq X_*)\}$ . For all  $\bar{\eta} \in Y$  we define  $u_{\bar{\eta}}(X_*) = \{m \in [1, k] \mid \forall n < \omega ([\bar{\eta} \upharpoonright m]_n \not\subseteq X_*)\}$ .
- (2) We say that the triple  $(Y_*, Y, X_*)$  is *f-closed* if  $(Y_*, Y)$  is  $\Lambda$ -closed,  $X_* \subseteq Y_*$  and, for all  $\bar{\eta} \in Y$ , either  $|u_{\bar{\eta}}(X_*)| \geq f$  or  $\bar{\eta} \in Y_{X_*}$ .
- (3) A regressive family  $\mathfrak{F}_{Y_*Y}$  of branch-like elements is called  $(Y, X_*)$ -*suitable* if  $[b_{\bar{\eta}}] \subseteq X_*$  for all  $\bar{\eta} \in Y_{X_*}$ .
- (4) Consider the map  $\rho_{Y_*X_*} : \bar{B}_{Y_*} \rightarrow \bar{B}_{Y_*}$  defined on the elements of  $Y_*$  by

$$e_{\bar{\nu}} \rho_{Y_*X_*} = \begin{cases} 0, & \text{if } \bar{\nu} \in X_* \\ e_{\bar{\nu}}, & \text{if } \bar{\nu} \in Y_* \setminus X_* \end{cases}$$

and extended by linearity. Define  $G_{Y_*YX_*} = \text{Im}(\rho_{Y_*X_*} \upharpoonright G_{Y_*Y})$ .

- OBSERVATION 41.
- (1) If  $(Y_*, Y)$  is  $\Lambda$ -closed,  $\bar{\eta} \in Y$  and  $X_* \subseteq Y_*$ , then  $u_{\bar{\eta}}(Y_*) \subseteq u_{\bar{\eta}}(X_*)$ . This fact will be used very often in the rest of this section.
  - (2) If  $Y \subseteq Z$ ,  $X_* \subseteq Y_* \subseteq Z_*$ , the pairs  $(Y_*, Y)$ ,  $(Z_*, Z)$  are  $\Lambda$ -closed and we have a regressive family  $\mathfrak{F}_{Z_*Z}$ , then  $G_{Y_*YX_*} \subseteq G_{Z_*ZX_*}$ .
  - (3) Moreover, if  $(Y_*, Y, X_*)$  and  $(Y'_*, Y', X'_*)$  are two *f-closed* triples such that  $Y'_* \subseteq Y_*$ ,  $Y' \subseteq Y$ ,  $X'_* \subseteq X_*$  and  $(Y'_* \setminus X'_*) \cap X_* = \emptyset$ , then  $\rho_{Y'_*X'_*} \subseteq \rho_{Y_*X_*}$ .

THEOREM 42. Let  $f \in [1, k]$ ,  $(Y_*, Y, X_*)$  be an  $f$ -closed triple and  $\mathfrak{F}_{Y_*Y}$  be regressive and  $(Y, X_*)$ -suitable. Put  $X = Y_{X_*}$ . Then  $(X_*, X)$  is  $\Lambda$ -closed and

$$G_{Y_*YX_*} \cong G_{Y_*Y}/G_{X_*X}$$

is  $\aleph_f$ -free, where  $G_{X_*X}$  is generated by  $\mathfrak{F}_{X_*X} = \mathfrak{F}_{Y_*X}$ .

PROOF. By the definition of  $Y_{X_*}$  there is some  $N_{\bar{\eta}} \leq N'_{\bar{\eta}} < \omega$  such that  $[\bar{\eta}]_{N'_{\bar{\eta}}} \subseteq X_*$ . Therefore  $(X_*, X)$  is  $\Lambda$ -closed. Furthermore, the  $(Y, X_*)$ -suitability of  $\mathfrak{F}_{Y_*Y}$  ensures that  $b_{\bar{\eta}} \in \overline{B}_{X_*}$  for all  $\bar{\eta} \in X$ . By Observation 38,  $G_{X_*X} \leq G_{Y_*Y}$ . We claim  $G_{X_*X} = \ker \rho_{Y_*X_*}$ . We have  $X_*\rho_{Y_*X_*} = 0$  by definition of  $\rho_{Y_*X_*}$ , and if  $\bar{\eta} \in X$ , then  $y_{\bar{\eta}N'_{\bar{\eta}}}\rho_{Y_*X_*} = 0$ . Moreover,  $[b_{\bar{\eta}}] \subseteq X_*$  since  $\mathfrak{F}_{Y_*Y}$  is  $(Y, X_*)$ -suitable. Hence,  $y'_{\bar{\eta}N'_{\bar{\eta}}}\rho_{Y_*X_*} = 0$ . Therefore,  $G_{X_*X} \subseteq \ker \rho_{Y_*X_*}$ . If  $g \in G_{Y_*Y} \cap \ker \rho_{Y_*X_*}$ , then

$$p^n g = \sum_{\bar{v} \in Y_*} a_{\bar{v}} e_{\bar{v}} + \sum_{\bar{\eta} \in Y \setminus X} a_{\bar{\eta}} y'_{\bar{\eta}N_{\bar{\eta}}} + \sum_{\bar{\eta} \in X} a_{\bar{\eta}} y'_{\bar{\eta}N'_{\bar{\eta}}}$$

for some  $n < \omega$ . Suppose that  $W = \{\bar{\eta} \in Y \setminus X \mid a_{\bar{\eta}} \neq 0\}$  is nonempty; and consider  $\bar{\eta} \in W$  of maximal norm. By the  $f$ -closedness of  $(Y_*, Y, X_*)$ , we obtain  $|u_{\bar{\eta}}(X_*)| \geq f$ . Take  $m \in u_{\bar{\eta}}(X_*)$  and  $n' < \omega$  large enough such that

$$\bar{\eta} \upharpoonright \langle m, n' \rangle \notin \left( \{\bar{v} \mid a_{\bar{v}} \neq 0\} \cup \bigcup_{\bar{\eta}' \in W \setminus \{\bar{\eta}\}} [y_{\bar{\eta}'}] \right) \cap X_*,$$

which are finitely many. It follows that  $a_{\bar{\eta}} y'_{\bar{\eta}N_{\bar{\eta}}}\rho_{Y_*X_*} \neq 0$ . Consequently,  $g \notin \ker \rho_{Y_*X_*}$ , which is a contradiction. Hence,  $W = \emptyset$  and  $a_{\bar{\eta}} = 0$  for all  $\bar{\eta} \in Y \setminus X$ . Similarly, it follows that  $a_{\bar{v}} = 0$  for all  $\bar{v} \in Y_* \setminus X_*$ . Therefore,  $g \in G_{X_*X}$ ; and our claim is proved, which implies that  $G_{Y_*YX_*} \cong G_{Y_*Y}/G_{X_*X}$ .

It is only left to show that  $G_{Y_*YX_*}$  is  $\aleph_f$ -free. We proceed as in the proof of Theorem 39 and construct a family  $\mathcal{C}$  of  $p$ -pure  $A$ -submodules of  $G_{Y_*YX_*}$  according to Definition 36. As in the proof of Theorem 39, for every  $H \in [G_{Y_*Y} \setminus G_{X_*X}]^{<\aleph_f}$ , we can find  $\Omega_* \subseteq Y_*$  and  $\Omega \subseteq Y$  such that  $|\Omega|, |\Omega_*| \leq |H| \cdot \aleph_0$  and

$$H \subseteq G_{\Omega_*\Omega} = \langle Ae_{\bar{\nu}}, Ay'_{\bar{\eta}N'_{\bar{\eta}}}, G_{X_*X} \mid \bar{\nu} \in \Omega_*, \bar{\eta} \in \Omega \rangle_* \subseteq G_{Y_*Y},$$

where, for all  $\bar{\eta} \in \Omega$ , we chose some  $N'_{\bar{\eta}} \geq N_{\bar{\eta}}$  such that  $[\bar{\eta}]_{N'_{\bar{\eta}}} \subseteq \Omega_*$ . Then  $G_{\Omega_*\Omega}/G_{X_*X}$  is  $<\aleph_f$ -generated. Let  $\Omega'_* = X_* \cup \bigcup_{\bar{\eta} \in \Omega} [\bar{\eta}]_{N'_{\bar{\eta}}} \cup [b_{\bar{\eta}N'_{\bar{\eta}}}]$  and  $\Delta = \Omega_* \setminus \Omega'_*$ . Notice that  $B_{\Delta}$  is a free summand of  $G_{\Omega_*\Omega}$  and that  $B_{\Delta}\rho_{Y_*X_*}$  is a free summand of  $G_{\Omega_*\Omega}\rho_{Y_*X_*}$ . Let

$$G_{\Omega} = \langle Ae_{\bar{\eta}\langle m,n \rangle}, Ae_{\bar{\nu}}, Ay'_{\bar{\eta}N'_{\bar{\eta}}}, G_{X_*X} \mid \bar{\eta} \in \Omega \setminus X, \bar{\nu} \in [b_{\bar{\eta}N'_{\bar{\eta}}}], m \in [1, k], n \in [N'_{\bar{\eta}}, \omega) \rangle_*.$$

Then,  $G_{\Omega'_*\Omega} \subseteq G_{\Omega}$  and

$$G_{\Omega}\rho_{Y_*X_*} \cong G_{\Omega}/G_{X_*X} \subseteq_* G_{Y_*Y}/G_{X_*X}.$$

Consequently,  $G_{\Omega}\rho_{Y_*X_*}$  is pure (see Fuchs [10], Vol.1, Lemma 26.1(ii)).

Suppose that  $|\Omega| = \aleph_{f-1}$ . Let  $F : \Lambda \rightarrow [\Lambda_*]^{<\aleph_0}$  be any regressive map such that  $\bar{\eta}F = [b_{\bar{\eta}}]$  for all  $\bar{\eta} \in \Omega$ . By taking  $u_{\bar{\eta}} = u_{\bar{\eta}}(X_*)$  for all  $\bar{\eta} \in \Omega \setminus X$  (which satisfy  $|u_{\bar{\eta}}(X_*)| \geq f$ ), we enumerate  $\Omega = \{\bar{\eta}^{\alpha} \mid \alpha < \zeta\}$  for some  $\zeta \in [\omega_{f-1}, \omega_f)$  according to the Freeness Proposition 11, and write

$$G_{\Omega} = \langle Ae_{\bar{\eta}^{\alpha}\langle m,n \rangle}, Ae_{\bar{\nu}}, Ae_{\bar{\nu}'}, Ay'_{\bar{\eta}^{\alpha}n}, Ay'_{\bar{\eta}'n'} \mid$$

$$\alpha < \zeta, m \in [1, k], n \in [N'_{\bar{\eta}}, \omega), \bar{\nu} \in [b_{\bar{\eta}^{\alpha}N'_{\bar{\eta}^{\alpha}}}], \bar{\nu}' \in X_*, \bar{\eta}' \in X, n' \in [N'_{\bar{\eta}'}, \omega) \rangle.$$

We also find  $\ell_\alpha \in u_{\bar{\eta}^\alpha}$  and  $n_\alpha < \omega$  such that

$$\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle \notin \{ \bar{\eta}^\beta \restriction \langle \ell_\alpha, n \rangle \mid \beta < \alpha \} \cup \bigcup \Omega_\alpha F$$

for all  $n \geq n_\alpha$ . Moreover,

$$\bar{\eta}^\alpha \restriction \langle \ell_\alpha, n \rangle \notin \{ \bar{\eta}^\beta \restriction \langle \ell_\alpha, n \rangle \mid \beta < \alpha \} \cup \bigcup \Omega_\alpha F \cup X_*$$

for infinitely many  $n \geq n_\alpha$ . Let

$$G_\alpha = \langle Ae_{\bar{\eta}^\beta \restriction \langle m, n \rangle}, Ae_{\bar{\nu}}, Ae_{\bar{\nu}'}, Ay'_{\bar{\eta}^\beta n}, Ay'_{\bar{\eta}' n'} \mid \\ \beta < \alpha, m \in [1, k], n \in [N'_{\bar{\eta}}, \omega), \bar{\nu} \in [b_{\bar{\eta}^\beta N'_{\bar{\eta}^\beta}}, \bar{\nu}' \in X_*, \bar{\eta}' \in X, n' \in [N_{\bar{\eta}'}, \omega) \rangle.$$

As in Theorem 39, it follows that  $G_0 = G_{X_* X}$ ,  $G_\zeta = G_\Omega$  and that  $G_\alpha$  is a direct summand of  $G_{\alpha+1}$ , with free complement in  $G_{\alpha+1}$ . We conclude that  $H\rho_{Y_* X_*}$  is a subset of the pure free  $A$ -submodule  $(B_\Delta \oplus G_\Omega)\rho_{Y_* X_*} = B_\Delta \rho_{Y_* X_*} \oplus G_\Omega \rho_{Y_* X_*}$  of  $G_{Y_* Y X_*}$ .

By setting  $\mathcal{C} = \{ (B_\Delta \oplus G_\Omega)\rho_{Y_* X_*} \mid |\Delta|, |\Omega| < \aleph_f \}$ , we conclude that  $G_{Y_* Y X_*}$  is  $\aleph_f$ -free.  $\square$

LEMMA 43. *Let  $(Z_*, Z, Y_*)$  and  $(Y_*, Y, X_*)$  be two  $f$ -closed triples such that  $Y = Z_{Y_*}$ .*

*Then*

(1)  *$(Z_*, Z, X_*)$  is  $f$ -closed and  $Z_{X_*} = Y_{X_*}$ .*

(2) *If  $\mathfrak{F}_{Z_* Z}$  is a regressive  $(Z, Y_*)$ -suitable family of branch-like elements such that*

*$\mathfrak{F}_{Z_* Y} \subseteq \mathfrak{F}_{Z_* Z}$  is  $(Y, X_*)$ -suitable, then*

$$G_{Z_* Z X_*} / G_{Y_* Y X_*} \cong G_{Z_* Z Y_*}.$$

PROOF. (1) On the one hand, it is immediate that  $Y_{X_*} \subseteq Z_{X_*}$ ; on the other hand,  $Z_{X_*} \subseteq Z_{Y_*} = Y$ , so  $Z_{X_*} \subseteq Y_{X_*}$ .

If  $\bar{\eta} \in Z \setminus Y$ , then  $|u_{\bar{\eta}}(Y_*)| \geq f$ , which implies that  $|u_{\bar{\eta}}(X_*)| \geq f$  because of  $u_{\bar{\eta}}(Y_*) \subseteq u_{\bar{\eta}}(X_*)$ . If  $\bar{\eta} \in Y$ , then either  $|u_{\bar{\eta}}(X_*)| \geq f$  or  $\bar{\eta} \in Y_{X_*}$  by the  $f$ -closedness of  $(Y_*, Y, X_*)$ .

(2) Observe that

$$G_{Z_*Z_{X_*}}/G_{Y_*Y_{X_*}} \cong (G_{Z_*Z}/G_{X_*Z_{X_*}}) / (G_{Y_*Y}/G_{X_*Y_{X_*}}) \cong G_{Z_*Z}/G_{Y_*Y} \cong G_{Z_*ZY_*}. \quad \square$$

DEFINITION 44. If  $(Y_*, Y)$  is  $\Lambda$ -closed and if  $\mathfrak{F}_{Y_*Y}$  is a regressive family of branch-like elements, then we say that  $\Omega_* \subseteq Y_*$  is  $\mathfrak{F}_{Y_*Y}$ -closed if  $(Y_*, Y, \Omega_*)$  is  $k$ -closed and  $\mathfrak{F}_{Y_*Y}$  is  $(Y, \Omega_*)$ -suitable.

DEFINITION 45. For  $Y \subseteq \Lambda$  and  $X_* \subseteq \Lambda_*$ , define

$$Y(X_*) = \{ \bar{\eta} \in Y \mid |u_{\bar{\eta}}(X_*)| < k \}.$$

THEOREM 46. If  $(Y_*, Y)$  is  $\Lambda$ -closed,  $X_* \subseteq Y_*$  and  $\mathfrak{F}_{Y_*Y}$  is a regressive family of branch-like elements, then there exists an  $\mathfrak{F}_{Y_*Y}$ -closed  $\Omega_* \subseteq Y_*$  such that  $X_* \subseteq \Omega_*$  and  $|\Omega_*| \leq |X_*|^{\aleph_0}$ .

PROOF. Let  $\Omega_0 = X_*$  and  $\Omega_\gamma = \bigcup_{\alpha < \gamma} \Omega_\alpha$  for all limit ordinals  $0 < \gamma < \omega_1$ . For  $\alpha < \omega_1$ , put

$$\Omega_{\alpha+1} = \Omega_\alpha \cup \bigcup_{\bar{\eta} \in Y(\Omega_\alpha)} [\bar{\eta}]_{N_{\bar{\eta}}} \cup [b_{\bar{\eta}}].$$



Observe that  $|\Omega_\alpha| \leq |X_*|^{\aleph_0}$  recursively, since  $|Y(\Omega_\alpha)| \leq |\Omega_\alpha|^{\aleph_0} \leq |X_*|^{\aleph_0}$ . By defining  $\Omega_* = \bigcup_{\alpha < \omega_1} \Omega_\alpha$ , we obtain  $|\Omega_*| \leq |X_*|^{\aleph_0}$  as well. If  $\bar{\eta} \in Y$  and  $|u_{\bar{\eta}}(\Omega_*)| < k$ , then  $\bar{\eta} \in Y(\Omega_*)$ , which in turn implies that  $\bar{\eta} \in Y(\Omega_\alpha)$  for some  $\alpha < \omega_1$ . It follows  $[\bar{\eta}]_{N_{\bar{\eta}}} \subseteq \Omega_{\alpha+1}$  and  $\bar{\eta} \in Y_{\Omega_{\alpha+1}} \subseteq Y_{\Omega_*}$ . Therefore,  $(Y_*, Y, \Omega_*)$  is  $k$ -closed. Now, if  $\bar{\eta} \in Y_{\Omega_*}$ , then  $\bar{\eta} \in Y(\Omega_\alpha)$  for some  $\alpha < \omega_1$ , which implies  $[b_{\bar{\eta}}] \subseteq \Omega_{\alpha+1} \subseteq \Omega_*$ . Hence,  $\mathfrak{F}_{Y_*Y}$  is  $(Y, \Omega_*)$ -suitable.  $\square$

OBSERVATION 47. In the previous theorem, it is necessary to take the union of the sets  $\Omega_\alpha$  according to  $\omega_1$  and not according to  $\omega$ , since it is possible that, for some  $\bar{\eta} \in Y$ ,  $|[\bar{\eta}] \cap \Omega_n| < \aleph_0$  for all  $n < \omega$ , but  $|[\bar{\eta}] \cap \Omega_\omega| = \aleph_0$ . Thus we cannot guarantee either that  $|u_{\bar{\eta}}(\Omega_\omega)| = k$  or  $\bar{\eta} \in Y_{\Omega_\omega}$ .

DEFINITION 48. If  $(Y_*, Y)$  is  $\Lambda$ -closed,  $X_* \subseteq Y_*$  and if  $\mathfrak{F}_{Y_*Y}$  is a regressive family of branch-like elements, then we say that  $\Omega_* \subseteq Y_*$  is the *closure of  $X_*$*  with respect to  $(Y_*, Y)$  and  $\mathfrak{F}_{Y_*Y}$  provided  $\Omega_*$  is constructed as in Theorem 46. We write  $\Omega_* = \overline{X}_*(Y_*, Y, \mathfrak{F}_{Y_*Y})$ .

In the next lemmas and Theorem 55, we want to "cover" small submodules of a triple module by small triple modules. The "small covers" are obtained by simple closure arguments (like elementary submodels) from the given large triple module.

LEMMA 49. *Let  $(Z_*, Z, Y_*)$  and  $(Y_*, Y, X_*)$  be two  $f$ -closed triples such that  $Y = Z_{Y_*}$ ,  $\mathfrak{F}_{Z_*Z}$  is regressive and  $(Z, Y_*)$ -suitable,  $\mathfrak{F}_{Z_*Y} \subseteq \mathfrak{F}_{Z_*Z}$  is  $(Y, X_*)$ -suitable and  $H \subseteq G_{Y_*YX_*}$ . Then there exists  $\Omega_* \subseteq Y_*$  such that*

- (1)  $|\Omega_*| \leq |H|^{\aleph_0}$ ,
- (2)  $(Z_*, Z, X_* \cup \Omega_*)$  and  $(X_* \cup \Omega_*, Y', X_*)$  are  $f$ -closed, where  $Y' = Z_{X_* \cup \Omega_*} = Y_{X_* \cup \Omega_*}$ ,
- (3)  $\mathfrak{F}_{Z_*Z}$  is  $(Z, X_* \cup \Omega_*)$ -suitable,
- (4)  $\mathfrak{F}_{Z_*Y'}$  is  $(Y', X_*)$ -suitable and
- (5)  $H \subseteq G_{X_* \cup \Omega_* Y' X_*} \subseteq G_{Y_* Y X_*} \subseteq G_{Z_* Z X_*}$ .

PROOF. (1) For every  $g \in H$ , choose some  $g' \in G_{Y_* Y}$  such that  $g' \rho_{Z_* X_*} = g$ ; and gather them in a set  $H'$ . Clearly  $[H] \subseteq [H'] \setminus X_*$ . Put  $\Omega_* = \overline{[H']}(Y_*, Y, \mathfrak{F}_{Z_* Y})$ . Then  $|\Omega_*| \leq |H|^{\aleph_0}$  follows by Theorem 46.

- (2) For  $\bar{\eta} \in Z$ , we have either  $|u_{\bar{\eta}}(Y_*)| \geq f$  or  $\bar{\eta} \in Z_{Y_*}$ . If  $|u_{\bar{\eta}}(Y_*)| \geq f$ , then  $|u_{\bar{\eta}}(X_* \cup \Omega_*)| \geq f$  since  $X_* \cup \Omega_* \subseteq Y_*$ . Otherwise, we have  $\bar{\eta} \in Z_{Y_*} = Y$ , so either  $|u_{\bar{\eta}}(X_*)| \geq f$  or  $\bar{\eta} \in Y_{X_*} \subseteq Z_{X_* \cup \Omega_*}$ . If  $|u_{\bar{\eta}}(X_*)| \geq f$ , then either  $|u_{\bar{\eta}}(\Omega_*)| = k$  or  $\bar{\eta} \in Y_{\Omega_*} \subseteq Z_{X_* \cup \Omega_*}$  since  $(Y_*, Y, \Omega_*)$  is  $k$ -closed. If  $|u_{\bar{\eta}}(\Omega_*)| = k$ , then  $|u_{\bar{\eta}}(X_* \cup \Omega_*)| = |u_{\bar{\eta}}(X_*)| \geq f$ . Hence,  $(Z_*, Z, X_* \cup \Omega_*)$  is  $f$ -closed.

Notice that  $Y_{X_* \cup \Omega_*} \subseteq Z_{X_* \cup \Omega_*}$  and that  $Z_{X_* \cup \Omega_*} \subseteq Z_{Y_*} = Y$ . Consequently,  $Z_{X_* \cup \Omega_*} \subseteq Y_{X_* \cup \Omega_*}$ . Since  $(Y_*, Y, X_*)$  is  $f$ -closed and  $Y'_{X_*} = Y_{X_*}$ ,  $(X_* \cup \Omega_*, Y', X_*)$  is also  $f$ -closed.

- (3) Let  $\bar{\eta} \in Y'$ . If  $|u_{\bar{\eta}}(\Omega_*)| = k$ , then  $\bar{\eta} \in Y_{X_*}$ , so  $[b_{\bar{\eta}}] \subseteq X_*$  because  $\mathfrak{F}_{Z_* Y}$  is  $(Y, X_*)$ -suitable. If  $\bar{\eta} \in Y_{\Omega_*}$ , then  $[b_{\bar{\eta}}] \subseteq \Omega_*$  by Theorem 46.
- (4) Follows from the  $(Y, X_*)$ -suitability of  $\mathfrak{F}_{Z_* Y}$  since  $Y'_{X_*} = Y_{X_*}$ .
- (5) The second inclusion follows from Observation 41. □

LEMMA 50. *Let  $(Z_*, Z, Y_*)$  and  $(Y_*, Y, X_*)$  be two  $f$ -closed triples such that  $Y = Z_{Y_*}$ ,  $\mathfrak{F}_{Z_*Z}$  is regressive and  $(Z, Y_*)$ -suitable,  $\mathfrak{F}_{Z_*Y} \subseteq \mathfrak{F}_{Z_*Z}$  is  $(Y, X_*)$ -suitable and  $K \subseteq G_{Z_*ZX_*}$ . Then there exist  $\Omega \subseteq Z$  and  $\Omega_* \subseteq Z_*$  such that*

- (1)  $|\Omega|, |\Omega_*| \leq |K| \cdot \aleph_0$ ,
- (2)  $(Y_* \cup \Omega_*, Y \cup \Omega, Y_*)$  is  $f$ -closed and  $(Y \cup \Omega)_{Y_*} = Y$ ,
- (3)  $\mathfrak{F}_{Z_*(Y \cup \Omega)}$  is  $(Y \cup \Omega, Y_*)$ -suitable,
- (4)  $K \subseteq G_{(Y_* \cup \Omega_*)(Y \cup \Omega)X_*} \subseteq G_{Z_*ZX_*}$  and
- (5) If  $(Z_*, Z, X_*)$  is  $f'$ -closed, then  $(Y_* \cup \Omega_*, Y \cup \Omega, X_*)$  is also  $f'$ -closed.

PROOF. (1) Let  $Z'_* = Z_* \setminus X_*$  and  $Z' = Z \setminus Z_{X_*}$ . For every  $g \in K$ , choose

some  $g' \in G_{Z'_*Z'}$  such that  $g'\rho_{Z_*X_*} = g$ ; and gather them in a set  $K'$ . Clearly

$[K] \subseteq [K'] \setminus X_*$ . Let  $\Omega_* = ([K'] \cup [[K']_\Lambda]) \cap Z_* \subseteq Z'_*$  and  $\Omega = Z_{\Omega_*}$ .

(2) By the  $\Lambda$ -closedness of  $(Y_*, Y)$ , it follows that  $Y = Y_{Y_*} \subseteq (Y \cup \Omega)_{Y_*} \subseteq Z_{Y_*} = Y$ .

Moreover,  $(Y_* \cup \Omega_*, Y \cup \Omega, Y_*)$  is  $f$ -closed, since  $(Z_*, Z, Y_*)$  is  $f$ -closed.

(3) Is immediate by the  $(Z, Y_*)$ -suitability of  $\mathfrak{F}_{Z_*Z}$  and the fact that  $(Y \cup \Omega)_{Y_*} = Z_{Y_*}$ .

(4) The second inclusion follows from Observation 41.

(5) Assume  $(Z_*, Z, X_*)$  is  $f'$ -closed; and let  $\bar{\eta} \in Y \cup \Omega$ . Then either  $|u_{\bar{\eta}}(X_*)| \geq f'$

or  $\bar{\eta} \in Z_{X_*} = Y_{X_*} \subseteq (Y \cup \Omega)_{X_*}$ . □

THEOREM 51. *Let  $(Z_*, Z, Y_*)$  and  $(Y_*, Y, X_*)$  be two  $f$ -closed triples such that  $Y = Z_{Y_*}$ ,  $\mathfrak{F}_{Z_*Z}$  is regressive and  $(Z, Y_*)$ -suitable,  $\mathfrak{F}_{Z_*Y} \subseteq \mathfrak{F}_{Z_*Z}$  is  $(Y, X_*)$ -suitable,  $H \subseteq G_{Y_*YX_*}$  and  $K \subseteq G_{Z_*ZX_*}$  such that  $|H|, |K| \leq \kappa$ . Then there exist  $Z'_*, Y'_*, X'_* \subseteq \Lambda_*$ ,  $Z', Y' \subseteq \Lambda$  such that*

- (1)  $Z'_* \subseteq Z_*$ ,  $Y'_* \subseteq Y_*$ ,  $X'_* \subseteq X_*$ ,  $Z' \subseteq Z \setminus Z_{X_*}$ ,  $Y' = Z'_{Y'_*} = Z_{Y'_*} \setminus Z_{X_*}$ ,
- (2)  $Z'_{X_*} = Y'_{X_*} = \emptyset$  and  $|Z'_*|, |Z'|, |Y'_*|, |Y'|, |X'_*| \leq \kappa^{\aleph_0}$ ,
- (3) the triples  $(Z'_*, Z', Y'_*)$ ,  $(Y'_*, Y', X'_*)$  are  $f$ -closed,
- (4)  $\mathfrak{F}_{Z_*Z'} \subseteq \mathfrak{F}_{Z_*Z}$  is  $(Z', Y'_*)$ -suitable and  $\mathfrak{F}_{Z_*Y'} \subseteq \mathfrak{F}_{Z_*Y}$  is  $(Y', X'_*)$ -suitable,
- (5)  $H \subseteq G_{Y'_*Y'X'_*} \subseteq G_{Y_*YX_*}$  and  $K \subseteq G_{Z'_*Z'X'_*} \subseteq G_{Z_*ZX_*}$ ,
- (6) If  $(Z_*, Z, X_*)$  is  $f'$ -closed, then  $(Z'_*, Z', X'_*)$  is also  $f'$ -closed.

PROOF. Apply Lemma 49 to  $H$  to obtain  $\Omega_*^1$ ; and let  $Y_1 = Z_{X_* \cup \Omega_*^1} = Y_{X_* \cup \Omega_*^1}$ . Apply Lemma 50 to  $K$  to obtain  $\Omega^2$  and  $\Omega_*^2$ . Define  $Z'_* = \Omega_*^1 \cup \Omega_*^2$ ,  $Y'_* = \Omega_*^1$ ,  $X'_* = X_* \cap \Omega_*^1$ ,  $Z' = (Y_1 \cup \Omega^2) \setminus (Y_1 \cup \Omega^2)_{X_*}$ . Then  $Z'_{Y'_*} = (Y_1 \cup \Omega^2)_{\Omega_*^1} \setminus (Y_1 \cup \Omega^2)_{X_*} = Z_{Y'_*} \setminus Z_{X_*}$ . Observe that  $\Omega^2 \setminus Z_{X_*} \subseteq Z'$  (which justifies condition (f) from Theorem 55).

(1) and (2) are immediate by definition.

- (3) Let  $\bar{\eta} \in Z'$ . If  $\bar{\eta} \in Y_1$ , then  $\bar{\eta} \in Z'_{Y'_*} \subseteq Z'_{Z'_*}$ . Otherwise,  $\bar{\eta} \in \Omega^2$ , and  $\bar{\eta} \in Z'_{\Omega_*^1 \cup \Omega_*^2} = Z'_{Z'_*}$ . Therefore,  $(Z'_*, Z')$  is  $\Lambda$ -closed. If  $|u_{\bar{\eta}}(X_* \cup Y'_*)| \geq f$ , then  $|u_{\bar{\eta}}(Y'_*)| \geq f$ . Otherwise,  $|u_{\bar{\eta}}(X_* \cup Y'_*)| < f$ , and  $\bar{\eta} \in (Y_1 \cup \Omega^2)_{X_* \cup Y'_*} \cap Z' = Z'_{Y'_*}$ . Therefore,  $(Z'_*, Z', Y'_*)$  is  $f$ -closed.

By definition of  $Y'$ , the pair  $(Y'_*, Y')$  is  $\Lambda$ -closed. Let  $\bar{\eta} \in Y' \subseteq Y_1$ . It is not possible to have  $|u_{\bar{\eta}}(X'_*)| < f$ , since it would imply  $|u_{\bar{\eta}}(X_*)| < f$ , so

$\bar{\eta} \in (Y_1)_{X_*} \cap Y' = \emptyset$ . Hence,  $|u_{\bar{\eta}}(X'_*)| \geq f$  for all  $\bar{\eta} \in Y'$  and  $(Y'_*, Y', X'_*)$  is  $f$ -closed.

(4) If  $\bar{\eta} \in Z'_{Y'_*} = Y'$ , then  $[b_{\bar{\eta}}] \subseteq Y'_* = \Omega_*^1$  by construction of  $\Omega_*^1$ , so  $\mathfrak{F}_{Z_*Z'} \subseteq \mathfrak{F}_{Z_*Z}$  is  $(Z', Y'_*)$ -suitable. Since  $Y'_{X'_*} = \emptyset$ ,  $\mathfrak{F}_{Z_*Y'} \subseteq \mathfrak{F}_{Z_*Y}$  is  $(Y', X'_*)$ -suitable.

(5) Follows by Observation 41.

(6) Let  $\bar{\eta} \in Z'$  and suppose  $|u_{\bar{\eta}}(X'_*)| < f'$ . It follows  $|u_{\bar{\eta}}(X_*)| < f'$ , so  $\bar{\eta} \in Z_{X_*} \cap Z' = \emptyset$ , which is a contradiction. Hence,  $|u_{\bar{\eta}}(X'_*)| \geq f'$  for all  $\bar{\eta} \in Z'$ ; and  $(Z'_*, Z', X'_*)$  is  $f'$ -closed.  $\square$

OBSERVATION 52. It is worth to remark that the construction of the triple  $(Y'_*, Y', X'_*)$  obtained in the last theorem depends only on  $(Y_*, Y, X_*)$  and  $H$ , and not on  $(Z_*, Z)$ .

DEFINITION 53. Let  $v = (\bar{\nu}_1, \bar{\nu}_2) \in \Lambda_* \times \Lambda_*$  and  $X_* \subseteq Y_* \subseteq \Lambda_*$ .

- (1)  $\bar{\nu}_1$  and  $\bar{\nu}_2$  are *comparable* if there exists a unique  $\bar{\eta}^v \in \Lambda$  such that  $\bar{\nu}_1, \bar{\nu}_2 \in [\bar{\eta}^v]$ .
- (2) Define  $\text{Cp}(X_*) = \{v = (\bar{\nu}_1, \bar{\nu}_2) \mid \bar{\nu}_1, \bar{\nu}_2 \in X_* \text{ are comparable}\} \subseteq X_* \times X_*$ .
- (3) We say that  $X_*$  is *pairwise closed for*  $Y_*$  if  $[\bar{\eta}^v] \cap Y_* \subseteq X_*$  for all  $v \in \text{Cp}(X_*)$ .

LEMMA 54. If  $X_* \subseteq Y_* \subseteq \Lambda_*$ , then there is a minimal pairwise closed set  $\text{PC}(X_*, Y_*)$  such that  $X_* \subseteq \text{PC}(X_*, Y_*) \subseteq Y_*$  and  $|\text{PC}(X_*, Y_*)| \leq |X_*| \cdot \aleph_0$ .

PROOF. Let  $X_*^0 = X_*$ . For  $n < \omega$ , let

$$X_*^{n+1} = X_*^n \cup \bigcup_{v \in \text{Cp}(X_*^n)} ([\bar{\eta}^v] \cap Y_*).$$

Put

$$\text{PC}(X_*, Y_*) = \bigcup_{n \in \omega} X_*^n.$$

Since  $|\text{Cp}(X_*)| \leq |X_*|$ , we have  $|\text{PC}(X_*, Y_*)| \leq |X_*| \cdot \aleph_0$ .  $\square$

**THEOREM 55.** *Let  $f \in [2, k]$ ,  $(Z_*, Z, Y_*)$  and  $(Y_*, Y, X_*)$  be two  $f$ -closed triples such that  $Y = Z_{Y_*}$  and  $\mathfrak{F}_{Z_*Z}$  be regressive and  $(Z, Y_*)$ -suitable such that  $\mathfrak{F}_{Z_*Y} \subseteq \mathfrak{F}_{Z_*Z}$  is  $(Y, X_*)$ -suitable. Let  $\Omega_*^1 \subseteq Y_*$ ,  $\Omega_*^2, \Omega_*^{2,n} \subseteq Z_*$  ( $n < \omega$ ),  $Z', Z_n \subseteq Z$  ( $n < \omega$ ) be such that*

- (a)  $\Omega_*^1$  is  $\mathfrak{F}_{Z_*Y}$ -closed,  $\Omega_*^2, \Omega_*^{2,n}$  are  $\mathfrak{F}_{Z_*Z}$ -closed,
- (b)  $\text{PC}(\Omega_*^{2,n}, Z_*) \subseteq \Omega_*^{2,n+1}$ ,
- (c)  $(\Omega_*^1 \cup \Omega_*^2, Z', \Omega_*^1)$ ,  $(\Omega_*^1 \cup \Omega_*^{2,n}, Z_n, \Omega_*^1)$  ( $n < \omega$ ) and  $(\Omega_*^1, Y', X_* \cap \Omega_*^1)$  are  $f$ -closed with  $Y' = Z'_{\Omega_*^1} = (Z_n)_{\Omega_*^1}$  ( $n < \omega$ ),
- (d)  $\mathfrak{F}_{Z_*Z'}$  is  $(Z', \Omega_*^1)$ -suitable,  $\mathfrak{F}_{Z_*Z_n}$  is  $(Z_n, \Omega_*^1)$ -suitable ( $n < \omega$ ) and  $\mathfrak{F}_{Z_*Y'}$  is  $(Y', X_* \cap \Omega_*^1)$ -suitable,
- (e)  $Z'_{X_*} = Y'_{X_*} = (Z_n)_{X_*} = \emptyset$  and
- (f)  $Z_{\Omega_*^2} \setminus Z_{X_*} \subseteq Z'$  and  $Z_{\Omega_*^{2,n}} \setminus Z_{X_*} \subseteq Z_n$ .

If we let

- (i)  $Z'' = \Omega_*^1 \cup \Omega_*^2 \cup \bigcup_{n < \omega} \Omega_*^{2,n}$ ,
- (ii)  $Y'' = \Omega_*^1 \cup \bigcup_{n < \omega} \Omega_*^{2,n}$ ,
- (iii)  $X'' = X_* \cap \Omega_*^1$ ,
- (iv)  $Z'' = Z' \cup \bigcup_{n < \omega} Z_n$  and
- (v)  $Y'' = Y' \cup \bigcup_{n < \omega} Z_n$ ,

then the following holds:

- (1)  $Z'' \subseteq Z \setminus Z_{X_*}$  and  $Z''_{X_*} = Y''_{X_*} = \emptyset$ ,
- (2)  $(Z'', Z'', Y'')$  is  $(f-1)$ -closed,  $(Y'', Y'', X'')$  is  $f$ -closed and  $Y'' = Z''_{Y''}$ ,
- (3)  $\mathfrak{F}_{Z_* Z''}$  is  $(Z'', Y'')$ -suitable and  $\mathfrak{F}_{Z_* Y''}$  is  $(Y'', X'')$ -suitable,
- (4)  $G_{Y'' Y'' X''} = G_{\Omega_*^1 Y' X_* \cap \Omega_*^1} + \sum_{n < \omega} G_{(\Omega_*^1 \cup \Omega_*^{2,n}) Z_n X_* \cap \Omega_*^1}$  and  
 $G_{Z'' Z'' X''} = G_{(\Omega_*^1 \cup \Omega_*^2) Z' X_* \cap \Omega_*^1} + \sum_{n < \omega} G_{(\Omega_*^1 \cup \Omega_*^{2,n}) Z_n X_* \cap \Omega_*^1}$ , and
- (5) if  $(Z_*, Z, X_*)$  is  $f'$ -closed then  $(Z'', Z'', X'')$  is also  $f'$ -closed.

PROOF. (1) It is immediate that  $Z', Z_n \subseteq Z \setminus Z_{X_*}$  by (e). The rest is clear.

- (2)  $(Z'', Z'')$  and  $(Y'', Y'')$  are  $\Lambda$ -closed since  $(\Omega_*^1 \cup \Omega_*^2, Z')$ ,  $(\Omega_*^1, Y')$  and all pairs  $(\Omega_*^1 \cup \Omega_*^{2,n}, Z_n)$  are  $\Lambda$ -closed.

Let  $\bar{\eta} \in Z''$ . If  $|u_{\bar{\eta}}(\Omega_*^1)| < f$ , then  $\bar{\eta} \in Z''_{\Omega_*^1} \subseteq Z''_{Y''}$  since  $(\Omega_*^1 \cup \Omega_*^2, Z', \Omega_*^1)$  and all triples  $(\Omega_*^1 \cup \Omega_*^{2,n}, Z_n, \Omega_*^1)$  are  $f$ -closed. Now assume that  $|u_{\bar{\eta}}(\Omega_*^1)| \geq f$ . On the one hand, if  $|u_{\bar{\eta}}(\Omega_*^1) \setminus u_{\bar{\eta}}(Y'')| > 1$ , then there exist distinct  $m_1, m_2 \in u_{\bar{\eta}}(\Omega_*^1) \setminus u_{\bar{\eta}}(Y'')$  such that  $[\bar{\eta} \upharpoonright m_1]_{n_1}, [\bar{\eta} \upharpoonright m_2]_{n_2} \subseteq Y''$  for some  $n_1, n_2 < \omega$ . Without loss of generality,  $\bar{\eta} \upharpoonright \langle m_1, n_1 \rangle, \bar{\eta} \upharpoonright \langle m_2, n_2 \rangle \in Y'' \setminus \Omega_*^1 \subseteq \bigcup_{n < \omega} \Omega_*^{2,n}$ . Thus,  $\bar{\eta} \upharpoonright \langle m_1, n_1 \rangle, \bar{\eta} \upharpoonright \langle m_2, n_2 \rangle \in \Omega_*^{2,n}$  for some  $n < \omega$ . Since  $(\bar{\eta} \upharpoonright \langle m_1, n_1 \rangle, \bar{\eta} \upharpoonright \langle m_2, n_2 \rangle) \in \text{Cp}(\Omega_*^{2,n})$  and  $\Omega_*^{2,n} \subseteq \text{PC}(\Omega_*^{2,n}, Z_*) \subseteq \Omega_*^{2,n+1}$ , it follows that  $[\bar{\eta}]_{N_{\bar{\eta}}} \subseteq [\bar{\eta}] \cap Z_* \subseteq \text{PC}(\Omega_*^{2,n}, Z_*) \subseteq \Omega_*^{2,n+1} \subseteq Y''$ . Hence,  $\bar{\eta} \in Z''_{Y''}$ . On the other hand, if  $|u_{\bar{\eta}}(\Omega_*^1) \setminus u_{\bar{\eta}}(Y'')| \leq 1$ , then  $|u_{\bar{\eta}}(Y'')| \geq f-1$ . Therefore  $(Z'', Z'', Y'')$  is  $(f-1)$ -closed.

If  $\bar{\eta} \in Y''$ , then  $|u_{\bar{\eta}}(X_*)| < f$  is not possible since that would imply  $|u_{\bar{\eta}}(X_*)| < f$ . By the  $f$ -closedness of  $(Z_*, Z, X_*)$ , we would get  $\bar{\eta} \in Z_{X_*} \cap Y'' = Y''_{X_*} = \emptyset$ . Hence,  $|u_{\bar{\eta}}(X_*)| \geq f$  for all  $\bar{\eta} \in Y''$ ; and  $(Y'', Y'', X_*)$  is  $f$ -closed.

Clearly  $Y'' \subseteq Z''_{Y''}$ . If  $\bar{\eta} \in Z''_{Y''}$ , then in particular  $\bar{\eta} \in Z'$  or  $\bar{\eta} \in Z_n$  for some  $n < \omega$ . If  $|u_{\bar{\eta}}(\Omega_*^1)| < f$ , then  $\bar{\eta} \in Z'_{\Omega_*^1} = (Z_n)_{\Omega_*^1} = Y' \subseteq Y''$  by the  $f$ -closedness of the triples  $(\Omega_*^1 \cup \Omega_*^2, Z', \Omega_*^1)$  and  $(\Omega_*^1 \cup \Omega_*^{2,n}, Z_n, \Omega_*^1)$  for all  $n < \omega$ . Otherwise,  $|u_{\bar{\eta}}(\Omega_*^1)| \geq f \geq 2$ . Hence, there exist  $m_1, m_2 \in [1, k]$  such that for all  $n < \omega$ ,  $[\bar{\eta} \upharpoonright m_1]_n, [\bar{\eta} \upharpoonright m_2]_n \notin \Omega_*^1$ . Then we can find  $n_1, n_2 < \omega$  such that  $\bar{\eta} \upharpoonright \langle m_1, n_1 \rangle, \bar{\eta} \upharpoonright \langle m_2, n_2 \rangle \in \bigcup_{n \in \omega} \Omega_*^{2,n}$ . A similar argument as above shows that  $[\bar{\eta}]_{N_{\bar{\eta}}} \subseteq \Omega_*^{2,n+1}$  for some  $n < \omega$ , showing that  $\bar{\eta} \in Z_{\Omega_*^{2,n+1}}$ . Observe that  $Z'' \subseteq Z \setminus Z_{X_*}$  by the definition of  $Z''$  and (e). It follows that  $\bar{\eta} \in Z_{\Omega_*^{2,n+1}} \setminus Z_{X_*} \subseteq Z_{n+1} \subseteq Y''$ .

- (3) Let  $\bar{\eta} \in Z''_{Y''} = Y''$ . If  $\bar{\eta} \in Y'$ , then  $[b_{\bar{\eta}}] \subseteq \Omega_*^1 \subseteq Y''$  since  $\mathfrak{F}_{Z_*Z'}$  is  $(Z', \Omega_*^1)$ -suitable and the families  $\mathfrak{F}_{Z_*Z_n}$  are  $(Z_n, \Omega_*^1)$ -suitable for all  $n < \omega$ . Now suppose  $\bar{\eta} \in Y'' \setminus Y'$ . Notice that  $|u_{\bar{\eta}}(\Omega_*^1)| \geq f$  since  $|u_{\bar{\eta}}(\Omega_*^1)| < f$  would imply that for some  $n \in \omega$ ,  $\bar{\eta} \in (Z_n)_{\Omega_*^1} = Y'$  by the  $f$ -closedness of the triple  $(\Omega_*^1 \cup \Omega_*^{2,n}, Z_n, \Omega_*^1)$ . As before,  $\bar{\eta} \in Z_{\Omega_*^{2,n+1}}$ , so  $[b_{\bar{\eta}}] \subseteq \Omega_*^{2,n+1} \subseteq Y''$  by the  $\mathfrak{F}_{Z_*Z}$ -closedness of  $\Omega_*^{2,n+1}$ . This shows that  $\mathfrak{F}_{Z_*Z''}$  is  $(Z'', Y'')$ -suitable.

Since  $Y''_{X_*} = \emptyset$ ,  $\mathfrak{F}_{Z_*Y''}$  is automatically  $(Y'', X_*)$ -suitable.

- (4) This follows from Observation 41(3).

- (5) Let  $\bar{\eta} \in Z''$  and suppose  $|u_{\bar{\eta}}(X_*)| < f'$ . Then  $|u_{\bar{\eta}}(X_*)| < f'$  and so it follows that  $\bar{\eta} \in Z_{X_*} \cap Z'' = Z''_{X_*} = \emptyset$ , which is a contradiction. Hence,  $|u_{\bar{\eta}}(X_*)| \geq f'$  for all  $\bar{\eta} \in Z''$  and  $(Z'', Z'', X_*)$  is  $f'$ -closed.  $\square$



### 3. The Step Lemma

DEFINITION 56. Let  $0 \leq \ell < m \leq k$  and  $\bar{\xi} \in \Lambda^{[\ell+1, m]}$ . Then

$$\Lambda^{\bar{\xi}} = \Lambda^{[1, \ell]} \wedge \{\bar{\xi}\},$$

$$\Lambda_*^{\bar{\xi}} = \Lambda_*^{[1, \ell]} \wedge \{\bar{\xi}\},$$

$$\Lambda^{\bar{\xi}*} = \Lambda^{[1, \ell]} \wedge [\bar{\xi}].$$

Furthermore, if  $\eta \in {}^{\omega}\lambda_\ell$  and  $W \subseteq [\eta]$ , let

$$\Lambda_W^{\bar{\xi}} = \Lambda^{[1, \ell-1]} \wedge W \wedge \{\bar{\xi}\},$$

$$\Lambda_\eta^{\bar{\xi}*} = \Lambda^{[1, \ell-1]} \wedge \{\eta\} \wedge [\bar{\xi}].$$

Notice that  $\Lambda_*^{\bar{\xi}} \dot{\cup} \Lambda^{\bar{\xi}*} = [\Lambda^{\bar{\xi}}]$  and  $\Lambda_*^{\bar{\xi}} \cap \Lambda^{\bar{\xi}*} = \emptyset$ . If  $n < \omega$ , we write  $\Lambda_{\eta|n}^{\bar{\xi}}$  for  $\Lambda_{\{\eta|n\}}^{\bar{\xi}}$ .

LEMMA 57. Let  $f \in [1, k)$ ,  $\bar{\xi} \in \Lambda^{[f+1, k]}$  and  $C_* \subseteq \Lambda_*$  be countable such that  $\Lambda(C_*) = \emptyset$  (see Definition 45). Then the triple  $(C_* \cup [\Lambda^{\bar{\xi}}], \Lambda^{\bar{\xi}}, C_* \cup \Lambda^{\bar{\xi}*})$  is  $f$ -closed.

PROOF. It is immediate that  $(C_* \cup [\Lambda^{\bar{\xi}}], \Lambda^{\bar{\xi}})$  is  $\Lambda$ -closed (with  $N_{\bar{\eta}} = 0$  for all  $\bar{\eta} \in \Lambda^{\bar{\xi}}$ ) and that  $[1, f] \subseteq u_{\bar{\eta}}(C_* \cup \Lambda^{\bar{\xi}*})$  because of the properties of  $C_*$  and the fact that  $\Lambda_*^{\bar{\xi}} \cap \Lambda^{\bar{\xi}*} = \emptyset$ .  $\square$

DEFINITION 58. (1) For  $\bar{\nu} \in \Lambda \cup \Lambda_*$  define  $\text{orco } \bar{\nu} = \bigcup_{m \in [1, k]} \text{Im } \nu_m$  to be the ordinal content of  $\bar{\nu}$ . Furthermore, if  $Y_* \subseteq \Lambda \cup \Lambda_*$ , then  $\text{orco } Y_* = \bigcup_{\bar{\nu} \in Y_*} \text{orco } \bar{\nu}$ .

- (2) If  $S, T \subseteq \lambda_k$  and  $\tau : S \rightarrow T$  is a bijection, then  $\tau$  extends canonically to a bijection  $\tau : {}^\omega \geq S \rightarrow {}^\omega \geq T$ . If  $\bar{\nu} \in \Lambda \cup \Lambda_*$  and  $\text{orco } \bar{\nu} \subseteq S$ , then we define  $\bar{\nu}\tau = (\nu_1\tau, \dots, \nu_k\tau)$ .
- (3) Given a finite sequence  $\langle \lambda'_1, \dots, \lambda'_k \rangle$  of nonempty subsets of  $T$ , consider

$$\Lambda' = {}^\omega \lambda'_1 \times \dots \times {}^\omega \lambda'_k,$$

$$\Lambda'_m = \Lambda'^{[1, m-1]} \wedge {}^\omega > \lambda'_m \wedge \Lambda'^{[m+1, k]},$$

$$\Lambda'_* = \bigcup_{m \in [1, k]} \Lambda'_m$$

and

$$B' = \bigoplus_{\bar{\nu} \in \Lambda'_*} A e_{\bar{\nu}},$$

defined in the same way as  $\Lambda$ ,  $\Lambda_m$ ,  $\Lambda_*$  and  $B$ . For  $X_* \subseteq \Lambda_*$  such that  $\text{orco } X_* \subseteq S$  and  $(\text{Im } \nu_m)\tau \subseteq \lambda'_m$  for all  $\nu \in X_*$  and  $m \in [1, k]$ , the bijection  $\tau : S \rightarrow T$  extends to an  $A$ -module monomorphism  $\tau : \bar{B}_{X_*} \rightarrow \bar{B}'$  called *shift-isomorphism* (onto its image).

- (4) For  $X_* \subseteq \Lambda_*$ , we say that the bijection  $\tau : S \rightarrow T$  is  $X_*$ -admissible if  $\text{orco } X_* \subseteq S$  and  $X_*\tau \subseteq \Lambda_*$ .
- (5) If  $\tau : S \rightarrow T$  is an  $X_*$ -admissible bijection, then  $\tau$  extends canonically to an  $A$ -module monomorphism  $\tau : \bar{B}_{X_*} \rightarrow \bar{B}$  called  $X_*$ -admissible *shift-isomorphism* (onto its image).

OBSERVATION 59. It is easy to verify that admissible shift-isomorphisms preserve all the notions we have defined so far, namely  $f$ -closedness, suitability, pairwise-closedness, etc.

STEP LEMMA 60. *Let the following be given:*

(1)  $f \in [0, k)$ ,  $\bar{\xi} \in \Lambda^{[f+1, k]}$  and a countable  $C_* \subseteq \Lambda_*$  such that  $\|C_*\| < 0\xi_k$  and

$$\Lambda(C_*) = \emptyset.$$

(2)  $(J_*, J, I_*) = (J_*(\bar{\xi}), J(\bar{\xi}), I_*(\bar{\xi})) = (C_* \cup [\Lambda^{\bar{\xi}}], \Lambda^{\bar{\xi}}, C_* \cup \Lambda^{\bar{\xi}*})$ .

(3)  $(V_*, V, U_*)$  an  $(f+1)$ -closed triple.

(4)  $\mathfrak{F}_{V_*V}$  regressive and  $(V, U_*)$ -suitable.

(5)  $\varphi : B_{I_*} \rightarrow G_{V_*VU_*}$  a homomorphism such that  $z\varphi \notin G_{V_*VU_*}$  for some  $z \in \overline{B}_{C_*}$ .

Then, for all  $\bar{\eta} \in J$ , we can choose an element  $b_{\bar{\eta}} \in \overline{B}_{C_*}$  such that the family  $\mathfrak{F}_{J_*J} = \{x_{\bar{\eta}} = b_{\bar{\eta}} + y_{\bar{\eta}} \mid \bar{\eta} \in J\}$  is regressive and  $G_{J_*J}$  satisfies the following condition:

If  $(Z_*, Z, Y_*)$  and  $(Y_*, Y, X_*)$  are two  $(f+1)$ -closed triples such that  $Z_{Y_*} = Y$ ,  $\mathfrak{F}_{Z_*Z} = \{y'_{\bar{\eta}} = b'_{\bar{\eta}} + y_{\bar{\eta}} \mid \bar{\eta} \in Z, b'_{\bar{\eta}} \in \overline{B}_{Z_*}\}$  is regressive and  $(Z, Y_*)$ -suitable,  $\mathfrak{F}_{Y_*Y} \subseteq \mathfrak{F}_{Z_*Z}$  is  $(Y, X_*)$ -suitable and  $\tau$  is a  $V_*$ -admissible bijection such that  $(V_*, V, U_*)\tau = (Y_*, Y, X_*)$  and  $\mathfrak{F}_{V_*V}\tau = \mathfrak{F}_{Y_*Y}$ , then  $\varphi\tau : B_{I_*} \rightarrow G_{Y_*YX_*}$  does not extend to a homomorphism from  $G_{J_*J}$  to  $G_{Z_*ZX_*}$ .

PROOF. We proceed by induction on  $f$ . If  $f = 0$ , then  $\bar{\xi} \in \Lambda$ ,  $J = \{\bar{\xi}\}$  and  $I_* = J_*$  since  $\Lambda_*^{\bar{\xi}} = \emptyset$ , namely  $(J_*, J, I_*) = (C_* \cup [\bar{\xi}], \{\bar{\xi}\}, C_* \cup [\bar{\xi}])$ . Consequently, we only need to define one element  $x_{\bar{\xi}}$  by choosing an element  $b_{\bar{\xi}}$ . We claim that it suffices to take  $b_{\bar{\xi}} = \varepsilon z$  for  $\varepsilon \in \{0, 1\}$ . Suppose that there exist two homomorphisms

$$\varphi^\varepsilon : \langle B_{J_*}, \varepsilon z + y_{\bar{\xi}} \rangle_* \rightarrow G_{V_*VU_*}$$

extending  $\varphi$ . On the one hand,  $(z + y_{\bar{\xi}})\varphi^1 - y_{\bar{\xi}}\varphi^0 \in G_{V_*VU_*}$ . On the other hand, by continuity of these homomorphisms,  $(z + y_{\bar{\xi}})\varphi^1 - y_{\bar{\xi}}\varphi^0 = z\varphi \notin G_{V_*VU_*}$ , which is a contradiction. We conveniently choose  $b_{\bar{\xi}}$  to satisfy  $x_{\bar{\xi}}\varphi \notin G_{V_*VU_*}$ . The construction of  $G_{J_*J}$  is finished. Moreover, since  $G_{V_*VU_*} = G_{V_*V \setminus V_{U_*}U_*}$ , this choice does not depend on  $\mathfrak{F}_{V_*VU_*}$ .

Assume towards a contradiction that there exist two 1-closed triples  $(Z_*, Z, Y_*)$  and  $(Y_*, Y, X_*)$  with  $Z_{Y_*} = Y$ ,  $\mathfrak{F}_{Z_*Z} = \{y'_{\bar{\eta}} = b'_{\bar{\eta}} + y_{\bar{\eta}} \mid \bar{\eta} \in Z, b'_{\bar{\eta}} \in \overline{B}_{Z_*}\}$  regressive and  $(Z, Y_*)$ -suitable,  $\mathfrak{F}_{Y_*Y} \subseteq \mathfrak{F}_{Z_*Z}$   $(Y, X_*)$ -suitable and a  $V_*$ -admissible bijection  $\tau$  such that  $(V_*, V, U_*)\tau = (Y_*, Y, X_*)$ ,  $\mathfrak{F}_{V_*V}\tau = \mathfrak{F}_{Y_*Y}$  and a homomorphism  $\psi : G_{J_*J} \rightarrow G_{Z_*ZX_*}$  extending  $\varphi\tau$ . Consider the unique extensions  $\widehat{\varphi} : \widehat{G}_{J_*J} \rightarrow \widehat{G}_{V_*VU_*}$  and  $\widehat{\tau} : \widehat{G}_{V_*VU_*} \rightarrow \widehat{G}_{Y_*YX_*}$ . Then  $\widehat{\varphi}\widehat{\tau} \upharpoonright G_{J_*J} = \psi$  and  $x_{\bar{\xi}}\psi \in G_{Z_*ZX_*} \cap \widehat{G}_{Y_*YX_*}$ . On the one hand,  $\widehat{G}_{Y_*YX_*}/G_{Y_*YX_*}$  is  $p$ -divisible. On the other hand,  $G_{Z_*ZX_*}/G_{Y_*YX_*} \cong G_{Z_*ZY_*}$  is  $\aleph_1$ -free by Theorem 42 and Lemma 43. It follows that  $x_{\bar{\xi}}\psi = x_{\bar{\xi}}\widehat{\varphi}\widehat{\tau} \in G_{Y_*YX_*}$  and  $x_{\bar{\xi}}\widehat{\varphi} \in G_{Y_*YX_*}\tau^{-1} = G_{V_*VU_*}$  contradicting the choice of  $b_{\bar{\xi}}$ . Therefore, no such  $\psi$  exists.

Now assume  $f > 0$  and that the statement is true for  $f - 1$ . Let

$$\lambda_f^o = \{ \alpha < \lambda_f \mid \text{cf}(\alpha) = \omega \},$$

$$\Gamma_\alpha = \{ \eta \in {}^{\omega^\uparrow}\lambda_f \mid \sup \eta = \alpha \}$$

and

$$\Gamma = \dot{\bigcup}_{\alpha \in \lambda_f^o} \Gamma_\alpha.$$

Observe that  $\Gamma = {}^\omega\lambda_f$ . We well-order each  $\Gamma_\alpha$  to induce a well-order in

$$\Gamma = \{ \eta^\alpha \mid \alpha < \lambda_f \}$$

by lexicographical extension of these orders. For all  $\alpha < \lambda_f$ , let  $\bar{\xi}^\alpha = \eta^\alpha \wedge \bar{\xi} \in \Lambda^{[f,k]}$ .

In this way,  $\Lambda^{\bar{\xi}} = \bigcup_{\alpha < \lambda_f} \Lambda^{\bar{\xi}^\alpha}$ . For  $\eta \in {}^\omega\lambda_f$  and  $n < \omega$ , let  $G_{\eta|n} = B_{\Lambda^{\bar{\xi}}_{[\eta|n]}}$ ; and define  $\mathcal{G} = \{ G_{\eta|n} \mid \eta \in {}^\omega\lambda_f, n < \omega \}$ . Notice that  $G_{\eta|(n+1)} = G_{\eta|n} \oplus B_{\Lambda^{\bar{\xi}}_{\eta|(n+1)}}$ .

Take  $H_\varphi = \text{Im } \varphi \subseteq G_{V_* V U_*}$  and then construct an  $(f+1)$ -closed triple  $(V', V', U') = (\Omega_*, V', U_* \cap \Omega_*)$  from  $H_\varphi$  and  $(V_*, V, U_*)$  (see Observation 52) as described in the proof of Theorem 51 and such that  $|V'_*|, |V'|, |U'_*| \leq \lambda_f$ ,  $H_\varphi \subseteq G_{V'_* V' U'_*} \subseteq G_{V_* V U_*}$ ,  $V'_{U'_*} = \emptyset$  and  $\mathfrak{F}_{V_* V'}$  is  $(V', U'_*)$ -suitable. Fix  $\Delta \subseteq \lambda_{f+1} \setminus \text{orco } V'_*$  such that  $|\Delta| = \lambda_f$ . Define a sequence  $\langle \lambda'_1, \lambda'_2, \dots, \lambda'_k \rangle$  of sets of ordinals in the following manner: for all  $m \in [1, k]$ , let

$$\lambda'_m = \begin{cases} \lambda_m, & \text{if } m \in [1, f]; \\ \Delta \dot{\cup} \text{orco } V'_*, & \text{if } m \in [f+1, k]. \end{cases}$$

Analogous to the definition of  $\Lambda$ ,  $\Lambda_m$ ,  $\Lambda_*$  and  $B$ , we define

$$\Lambda' = {}^\omega\lambda'_1 \times \dots \times {}^\omega\lambda'_k,$$

$$\Lambda'_m = \Lambda'^{[1, m-1]} \wedge {}^\omega\lambda'_m \wedge \Lambda'^{[m+1, k]},$$

$$\Lambda'_* = \bigcup_{m \in [1, k]} \Lambda'_m,$$

$$B' = \bigoplus_{\bar{v} \in \Lambda'_*} A e_{\bar{v}}.$$

Let  $\mathcal{H} = \{ H \leq \overline{B'} \mid |H| \leq \lambda_{f-1} \}$  and consider the set  $\Theta$  of all 6-tuples  $(G, H, P, Q, R, \psi)$  where  $G \in \mathcal{G}$ ,  $H \in \mathcal{H}$ ,  $P \in [\Lambda'_*]^{\leq \lambda_{f-1}}$ ,  $Q \in [\Lambda']^{\leq \lambda_{f-1}}$ ,  $R \in [\overline{B'}]^{\leq \lambda_{f-1}}$  and  $\psi : G \rightarrow H$  is a homomorphism. Recall that  $\lambda_0 = |A|$  (as agreed on page 21) and notice that

$$|\Theta| = |\mathcal{G}| \cdot |\mathcal{H}| \cdot |\Lambda'_*|^{\lambda_{f-1}} \cdot |\Lambda'|^{\lambda_{f-1}} \cdot |\overline{B'}|^{\lambda_{f-1}} \cdot \lambda_{f-1}^{\lambda_{f-1}} = \lambda_f^{\lambda_{f-1}} = \lambda_f.$$

Take  $\overline{\lambda} = \langle \lambda_f \rangle$  and  $\overline{C} = \langle \Theta \rangle$  and apply the  $\overline{\lambda}$ -Black Box 15 to obtain a family of traps  $\langle g_\eta \mid \eta \in \Gamma \rangle = \langle g_{\eta^\alpha} \mid \alpha < \lambda_f \rangle$  with  $g_\eta : [\eta] \rightarrow \Theta$ . For every  $\alpha < \lambda_f$ , we will choose  $b_{\overline{\eta}}$  for all  $\overline{\eta} \in \Lambda^{\overline{\xi}^\alpha}$  by examining the trap  $g_{\eta^\alpha} : [\eta^\alpha] \rightarrow \Theta$  on its values  $(\eta^\alpha \upharpoonright n)g_{\eta^\alpha} = (G_{\alpha n}, H_{\alpha n}, P_{\alpha n}, Q_{\alpha n}, R_{\alpha n}, \psi_{\alpha n})$ . We will choose these elements by means of the induction hypothesis only under very specific conditions. Otherwise we will take  $b_{\overline{\eta}} = 0$  for all  $\overline{\eta} \in \Lambda^{\overline{\xi}^\alpha}$ . These conditions are:

If there exist  $Y, Z \subseteq \Lambda$ ,  $X_*, Y_*, Z_* \subseteq \Lambda_*$ ,  $\mathfrak{F}_{Z_*Z}$ ,  $\Omega_*^1 \subseteq Y_*$ ,  $Y' \subseteq Y$ , families of subsets  $\{ \Omega_*^{2,n} \subseteq Z_* \mid n < \omega \}$  and  $\{ Z_n \subseteq Z \mid n < \omega \}$ ,  $\tau$  and  $\sigma$  such that

- (1)  $(Z_*, Z, Y_*)$  and  $(Y_*, Y, X_*)$  are  $(f+1)$ -closed with  $Z_{Y_*} = Y$ ,
- (2)  $\mathfrak{F}_{Z_*Z}$  is a regressive  $(Z, Y_*)$ -suitable family of branch-like elements and  $\mathfrak{F}_{Y_*Y} \subseteq \mathfrak{F}_{Z_*Z}$  is  $(Y, X_*)$ -suitable,
- (3)  $\Omega_*^1$  is  $\mathfrak{F}_{Y_*Y}$ -closed and  $\Omega_*^{2,n}$  is  $\mathfrak{F}_{Z_*Z}$ -closed for all  $n < \omega$ ,
- (4)  $\text{PC}(\Omega_*^{2,n}, Z_*) \subseteq \Omega_*^{2,n+1}$ ,
- (5)  $(\Omega_*^1, Y', X_* \cap \Omega_*^1)$  and  $(\Omega_*^1 \cup \Omega_*^{2,n}, Z_n, \Omega_*^1)$  are  $(f+1)$ -closed with  $(Z_n)_{\Omega_*^1} = Y'$  for all  $n < \omega$ ,
- (6)  $\mathfrak{F}_{Z_*Y'}$  is  $(Y', X_* \cap \Omega_*^1)$ -suitable and  $\mathfrak{F}_{Z_*Z_n}$  is  $(Z_n, \Omega_*^1)$ -suitable for all  $n < \omega$ ,
- (7)  $G_{\Omega_*^1 Y' X_* \cap \Omega_*^1} \subseteq G_{Y_* Y X_*}$  and  $G_{\Omega_*^1 \cup \Omega_*^{2,n} Z_n X_* \cap \Omega_*^1} \subseteq G_{Z_* Z X_*}$  for all  $n < \omega$ ,
- (8)  $Y'_{X_*} = (Z_n)_{X_*} = \emptyset$  for all  $n < \omega$ ,

(9)  $Y' = Z_{\Omega_*^1} \setminus Z_{X_*}$  and  $Z_{\Omega_*^{2,n}} \setminus Z_{X_*} \subseteq Z_n \subseteq Z \setminus Z_{X_*}$  for all  $n < \omega$ ,

(10)  $\tau$  is a  $V_*$ -admissible bijection such that

$$(V_*, V, U_*)\tau = (Y_*, Y, X_*),$$

$$(\Omega_*, V', U_* \cap \Omega_*)\tau = (\Omega_*^1, Y', X_* \cap \Omega_*^1),$$

$$\mathfrak{F}_{V_*V}\tau = \mathfrak{F}_{Y_*Y} \text{ and } \mathfrak{F}_{V_*V'}\tau = \mathfrak{F}_{Y_*Y'},$$

(11)  $\sigma : \text{orco}(\Omega_*^1 \cup \bigcup_{n < \omega} \Omega_*^{2,n}) \rightarrow \Delta \cup \text{orco } V'_*$  is a shift-monomorphism such that

$$\sigma \upharpoonright \text{orco}(\Omega_*^1) = \tau^{-1} \upharpoonright \text{orco}(\Omega_*^1),$$

(12)  $G_{\alpha n} = G_{\eta^\alpha \upharpoonright n}$ ,  $H_{\alpha n} \subseteq G_{\Omega_*^1 \cup \Omega_*^{2,n} Z_n X_* \cap \Omega_*^1} \sigma$ ,  $P_{\alpha n} = \Omega_*^{2,n} \sigma$ ,  $Q_{\alpha n} = (Z_n \setminus Y')\sigma$ ,

$$R_{\alpha n} = \mathfrak{F}_{Z_* Z_n \setminus Y'} \sigma \text{ and}$$

(13) the maps  $\psi_{\alpha n} : G_{\alpha n} \rightarrow H_{\alpha n}$  extend each other.

Suppose that at some stage  $\alpha < \lambda_f$ , we find such  $(f+1)$ -closed triples  $(Z_{\alpha*}, Z_\alpha, Y_{\alpha*})$  and  $(Y_{\alpha*}, Y_\alpha, X_{\alpha*})$ , a regressive family  $\mathfrak{F}_{Z_{\alpha*}Z_\alpha}$  of branch-like elements,  $\Omega_{\alpha*}^1 \subseteq Y_{\alpha*}$ ,  $Y'_\alpha \subseteq Y_\alpha$ , families of subsets  $\{\Omega_{\alpha*}^{2,n} \subseteq Z_{\alpha*} \mid n < \omega\}$  and  $\{Z_{\alpha n} \subseteq Z_\alpha \mid n < \omega\}$  and bijections  $\tau_\alpha$  and  $\sigma_\alpha$  satisfying these conditions. Define  $G_\alpha = \bigcup_{n < \omega} G_{\alpha n} = B_{\Lambda_{[\eta^\alpha]}^\xi}$  and  $\psi_\alpha = \bigcup_{n < \omega} \psi_{\alpha n} : G_\alpha \rightarrow \sum_{n < \omega} H_{\alpha n}$ , so

$$\psi_\alpha \sigma_\alpha^{-1} : G_\alpha \rightarrow \sum_{n < \omega} G_{\Omega_{\alpha*}^1 \cup \Omega_{\alpha*}^{2,n} Z_{\alpha n} X_{\alpha*} \cap \Omega_{\alpha*}^1}.$$

According to Theorem 55 we define an  $(f+1)$ -closed triple  $(Y''_{\alpha*}, Y''_\alpha, X''_{\alpha*})$  as follows:

$$Y''_{\alpha*} = \Omega_{\alpha*}^1 \cup \bigcup_{n < \omega} \Omega_{\alpha*}^{2,n},$$

$$Y''_\alpha = Y'_\alpha \cup \bigcup_{n < \omega} Z_{\alpha n},$$

$$X''_{\alpha*} = X_{\alpha*} \cap \Omega_{\alpha*}^1.$$

Consider the triple  $(J_*(\bar{\xi}^\alpha), J(\bar{\xi}^\alpha), I_*(\bar{\xi}^\alpha)) = (C_* \cup [\Lambda^{\bar{\xi}^\alpha}], \Lambda^{\bar{\xi}^\alpha}, C_* \cup \Lambda^{\bar{\xi}^\alpha*})$ , and notice that

$$B_{I_*(\bar{\xi}^\alpha)} = B_{C_* \cup \Lambda^{\bar{\xi}^\alpha*}} = B_{C_* \cup \Lambda_{\eta^\alpha}^{\bar{\xi}^\alpha}} \oplus B_{\Lambda_{[\eta^\alpha]}^{\bar{\xi}^\alpha}} \subseteq B_{I_*(\bar{\xi})} \oplus B_{\Lambda_{[\eta^\alpha]}^{\bar{\xi}}} = B_{I_*(\bar{\xi})} \oplus G_\alpha.$$

Let  $\varphi'_\alpha = \left( \varphi \upharpoonright B_{C_* \cup \Lambda_{\eta^\alpha}^{\bar{\xi}^\alpha}} \right) \tau_\alpha \oplus \psi_\alpha \sigma_\alpha^{-1}$ . Then by definition,

$$\varphi'_\alpha : B_{I_*(\bar{\xi}^\alpha)} \rightarrow G_{\Omega_{\alpha*}^1 Y'_\alpha X_{\alpha*} \cap \Omega_{\alpha*}^1} + \sum_{n < \omega} G_{\Omega_{\alpha*}^1 \cup \Omega_{\alpha*}^{2,n} Z_{\alpha n} X_{\alpha*} \cap \Omega_{\alpha*}^1} = G_{Y''_{\alpha*} Y''_\alpha X''_{\alpha*}}.$$

We claim that  $G_{Y''_{\alpha*} Y''_\alpha X''_{\alpha*}} \cap \widehat{G}_{\Omega_{\alpha*}^1 Y'_\alpha X_{\alpha*} \cap \Omega_{\alpha*}^1} = G_{\Omega_{\alpha*}^1 Y'_\alpha X_{\alpha*} \cap \Omega_{\alpha*}^1}$ . On the one hand, let  $g \in G_{Y''_{\alpha*} Y''_\alpha X''_{\alpha*}} \cap \widehat{G}_{\Omega_{\alpha*}^1 Y'_\alpha X_{\alpha*} \cap \Omega_{\alpha*}^1}$ , and choose some preimage  $g' \in G_{Y''_{\alpha*} Y''_\alpha}$  such that  $g' \rho_{Y''_{\alpha*} X''_{\alpha*}} = g$ . Then we can write

$$g' = b + \sum_{\bar{\eta} \in Y''_\alpha} a_{\bar{\eta}} y'_{\bar{\eta}}$$

for some  $b \in B_{Y''_{\alpha*}}$  and  $a_{\bar{\eta}} \in A$  for all  $\bar{\eta} \in Y''_\alpha$ , where  $a_{\bar{\eta}} \neq 0$  only for finitely many elements  $\bar{\eta}$ . Also notice that  $[g'] \subseteq \Omega_{\alpha*}^1$  and  $Y''_\alpha \setminus (Y''_\alpha)_{X''_{\alpha*}} = Y''_\alpha$  since  $(Y''_\alpha)_{X''_{\alpha*}} = \emptyset$  by condition (8). Without loss of generality, we can take

$$g' = b + \sum_{\bar{\eta} \in Y''_\alpha \setminus Y'_\alpha} a_{\bar{\eta}} y'_{\bar{\eta}}$$

(otherwise just reduce  $g$  and  $g'$ ). Take  $\bar{\eta} \in [g']_\Lambda \subseteq Y''_\alpha \setminus Y'_\alpha$  of maximal norm. Since  $[g'] \subseteq \Omega_{\alpha*}^1$ , it follows by condition (9) that

$$\bar{\eta} \in (Y''_\alpha)_{\Omega_{\alpha*}^1} \setminus Y'_\alpha \subseteq (Z_\alpha \setminus (Z_\alpha)_{X_{\alpha*}})_{\Omega_{\alpha*}^1} \setminus Y'_\alpha \subseteq ((Z_\alpha)_{\Omega_{\alpha*}^1} \setminus (Z_\alpha)_{X_{\alpha*}}) \setminus Y'_\alpha = Y'_\alpha \setminus Y'_\alpha = \emptyset.$$

Therefore,  $g' = b \in G_{\Omega_{\alpha*}^1 Y'_\alpha}$  and  $g = g' \rho_{Y''_{\alpha*} X''_{\alpha*}} \in G_{\Omega_{\alpha*}^1 Y'_\alpha X_{\alpha*} \cap \Omega_{\alpha*}^1}$  showing

$$G_{Y''_{\alpha*} Y''_\alpha X''_{\alpha*}} \cap \widehat{G}_{\Omega_{\alpha*}^1 Y'_\alpha X_{\alpha*} \cap \Omega_{\alpha*}^1} \subseteq G_{\Omega_{\alpha*}^1 Y'_\alpha X_{\alpha*} \cap \Omega_{\alpha*}^1}.$$



On the other hand, it is immediate that  $G_{\Omega_{\alpha*}^1 Y_{\alpha}' X_{\alpha*} \cap \Omega_{\alpha*}^1} \subseteq G_{Y_{\alpha*}'' Y_{\alpha}'' X_{\alpha*}''} \cap \widehat{G}_{\Omega_{\alpha*}^1 Y_{\alpha}' X_{\alpha*} \cap \Omega_{\alpha*}^1}$ . This proves our claim. Furthermore,  $z\varphi'_{\alpha} = z\varphi\tau_{\alpha} \notin G_{Y_{\alpha*}'' Y_{\alpha}'' X_{\alpha*}''}$  since  $z\varphi'_{\alpha} \in G_{Y_{\alpha*}'' Y_{\alpha}'' X_{\alpha*}''}$  would imply  $z\varphi'_{\alpha} \in G_{\Omega_{\alpha*}^1 Y_{\alpha}' X_{\alpha*} \cap \Omega_{\alpha*}^1}$  by our previous claim. Hence,  $z\varphi'_{\alpha}\tau_{\alpha}^{-1} = z\varphi \in G_{\Omega_{*} V' U_{*} \cap \Omega_{*}} \subseteq G_{V_{*} V U_{*}}$ , which is a contradiction. In particular, since  $(Y_{\alpha*}'', Y_{\alpha}'', X_{\alpha*}'')$  is  $f$ -closed and  $\bar{\xi}^{\alpha} \in \Lambda^{[f,k]}$ , we can apply the induction hypothesis to choose  $b_{\bar{\eta}}$  for all  $\bar{\eta} \in \Lambda^{\bar{\xi}^{\alpha}}$ . This finishes the construction of  $G_{J_* J}$ .

Assume towards a contradiction that there exist two  $(f+1)$ -closed triples  $(Z_*, Z, Y_*)$  and  $(Y_*, Y, X_*)$  with  $Z_{Y_*} = Y$ , a regressive  $(Z, Y_*)$ -suitable family  $\mathfrak{F}_{Z_* Z}$  of branch-like elements such that  $\mathfrak{F}_{Y_* Y} \subseteq \mathfrak{F}_{Z_* Z}$  is  $(Y, X_*)$ -suitable, a  $V_*$ -admissible bijection  $\tau$  such that  $(V_*, V, U_*)\tau = (Y_*, Y, X_*)$ ,  $\mathfrak{F}_{V_* V}\tau = \mathfrak{F}_{Y_* Y}$  and a homomorphism  $\psi : G_{J_* J} \rightarrow G_{Z_* Z X_*}$  extending  $\varphi\tau$ .

Take  $H = \text{Im } \varphi\tau \subseteq G_{Y_* Y X_*}$  and  $K = \text{Im } \psi \subseteq G_{Z_* Z X_*}$ . Apply Theorem 51 to construct the sets  $\Omega_*^1, Y_1, \Omega_*^2, \Omega^2, Y'$  and  $Z'$  and the two  $(f+1)$ -closed triples  $(\Omega_*^1 \cup \Omega_*^2, Z', \Omega_*^1)$  and  $(\Omega_*^1, Y', X_* \cap \Omega_*^1)$  such that  $H \subseteq G_{\Omega_*^1 Y' X_* \cap \Omega_*^1} \subseteq G_{\Omega_*^1 \cup \Omega_*^2 Z' \Omega_*^1}$  and  $K \subseteq G_{\Omega_*^1 \cup \Omega_*^2 Z' \Omega_*^1} \subseteq G_{Z_* Z X_*}$ . Fix an injection  $\sigma : \text{orco}(\Omega_*^1 \cup \Omega_*^2) \rightarrow \Delta \cup \text{orco } V'_*$  such that  $\sigma \upharpoonright \text{orco}(\Omega_*^1) = \tau^{-1} \upharpoonright \text{orco}(\Omega_*^1)$ .

For all  $\eta \in {}^{\omega}\lambda_f$  and  $n < \omega$ , let  $K_{\eta \upharpoonright n} = \text{Im}(\psi \upharpoonright G_{\eta \upharpoonright n})$ . Put  $\Omega_*^{\eta \upharpoonright 0} = \emptyset$ ; and assume we have constructed  $\Omega_*^{\eta \upharpoonright n}$ . Let  $\Omega_*^{\eta \upharpoonright n} = Z_{\Omega_*^{\eta \upharpoonright n}}$  and  $Z^{\eta \upharpoonright n} = (Y_1 \cup \Omega_*^{\eta \upharpoonright n}) \setminus (Y_1 \cup \Omega_*^{\eta \upharpoonright n})_{X_*}$ . For every  $\bar{\nu} \in \Lambda_{\eta \upharpoonright n}^{\bar{\xi}}$ , choose some  $h_{\bar{\nu}} \in G_{Z_* Z}$  such that  $h_{\bar{\nu}}\rho_{Z_* X_*} = e_{\bar{\nu}}\psi$ . Apply Lemma 50 and Theorem 51 to find an  $\mathfrak{F}_{Z_* Z}$ -closed  $\Omega_*^{\eta \upharpoonright (n+1)}$  such that

$$\text{PC}(\Omega_*^{\eta \upharpoonright n}, Z_*) \cup \bigcup_{\bar{\nu} \in \Lambda_{\eta \upharpoonright n}^{\bar{\xi}}} [h_{\bar{\nu}}] \subseteq \Omega_*^{\eta \upharpoonright (n+1)}.$$

In this way, we obtain families  $\{ \Omega_*^{\eta \upharpoonright n} \subseteq Z_* \mid n < \omega \}$  and  $\{ Z^{\eta \upharpoonright n} \subseteq Z \mid n < \omega \}$  for all  $\eta \in {}^{\omega \uparrow} \lambda_f$  simultaneously.

Let  $\mathbf{g} : {}^{\omega \uparrow} \lambda_f \rightarrow \Theta$  be given by

$$\nu \mathbf{g} = (G^\nu, H^\nu, P^\nu, Q^\nu, R^\nu, \psi^\nu) = (G_\nu, K_\nu \sigma, \Omega_*^\nu \sigma, (Z^\nu \setminus Y') \sigma, \mathfrak{F}_{Z_* Z^\nu \setminus Y'} \sigma, (\psi \upharpoonright G_\nu) \sigma).$$

By the  $\bar{\lambda}$ -Black Box 15 there is some  $\eta^\alpha \in \Gamma$  such that  $g_{\eta^\alpha} \subseteq \mathbf{g}$ . It follows that

$$\begin{aligned} (\eta^\alpha \upharpoonright n) \mathbf{g} &= (G_{\alpha n}, H_{\alpha n}, P_{\alpha n}, Q_{\alpha n}, R_{\alpha n}, \psi_{\alpha n}) \\ &= (G_{\eta^\alpha \upharpoonright n}, K_{\eta^\alpha \upharpoonright n} \sigma, \Omega_*^{\eta^\alpha \upharpoonright n} \sigma, (Z^{\eta^\alpha \upharpoonright n} \setminus Y') \sigma, \mathfrak{F}_{Z_* Z^{\eta^\alpha \upharpoonright n} \setminus Y'} \sigma, (\psi \upharpoonright G_{\eta^\alpha \upharpoonright n}) \sigma). \end{aligned}$$

Consequently,  $Y, Z \subseteq \Lambda$ ,  $X_*, Y_*, Z_* \subseteq \Lambda_*$ ,  $\Omega_*^1 \subseteq Y_*$ ,  $Y' \subseteq Y$ ,  $\mathfrak{F}_{Z_* Z}$ ,  $\tau$ ,  $\sigma$  and the families  $\{ \Omega_*^{\eta^\alpha \upharpoonright n} \subseteq Z_* \mid n < \omega \}$  and  $\{ Z^{\eta^\alpha \upharpoonright n} \subseteq Z \mid n < \omega \}$  satisfy the specific conditions (1)-(13) of the construction. Let

$$Z'' = \Omega_*^1 \cup \Omega_*^2 \cup \bigcup_{n < \omega} \Omega_*^{\eta^\alpha \upharpoonright n},$$

$$Y'' = \Omega_*^1 \cup \bigcup_{n < \omega} \Omega_*^{\eta^\alpha \upharpoonright n},$$

$$X'' = X_* \cap \Omega_*^1,$$

$$Z'' = Z' \cup \bigcup_{n < \omega} Z^{\eta^\alpha \upharpoonright n} \text{ and}$$

$$Y'' = Y' \cup \bigcup_{n < \omega} Z^{\eta^\alpha \upharpoonright n}.$$

By Theorem 55,  $(Z'', Z'', Y'')$  is  $f$ -closed and  $(Y'', Y'', X'')$  is  $(f+1)$ -closed. Let  $\psi_\alpha = \bigcup_{n < \omega} \psi_{\alpha n}$  and

$$\varphi' = \left( \varphi \upharpoonright B_{C_* \cup \Lambda_{\eta^\alpha}^{\bar{\xi}_*}} \right) \tau \oplus \psi_\alpha \sigma^{-1} \subseteq \varphi \tau \oplus (\psi \upharpoonright G_\alpha) \subseteq \psi.$$

Since in the step  $\alpha$  of the construction of  $G_{J_* J}$  the conditions (1)-(13) for a serious choice of the  $b_{\bar{\eta}}$ 's were in fact satisfiable, we used some  $(f+1)$ -closed triples  $(Z_{\alpha*}, Z_\alpha, Y_{\alpha*})$  and

$(Y_{\alpha*}, Y_\alpha, X_{\alpha*})$ , a regressive family  $\mathfrak{F}_{Z_{\alpha*}Z_\alpha}$  of branch-like elements,  $\Omega_{\alpha*}^1 \subseteq Y_{\alpha*}$ ,  $Y'_\alpha \subseteq Y_\alpha$ , families  $\{\Omega_{\alpha*}^{2,n} \subseteq Z_{\alpha*} \mid n < \omega\}$  and  $\{Z_{\alpha,n} \subseteq Z_\alpha \mid n < \omega\}$ ,  $\tau_\alpha$  and  $\sigma_\alpha$  in this stage of the construction. Also from this stage, we get an  $(f+1)$ -closed triple  $(Y''_{\alpha*}, Y''_\alpha, X''_{\alpha*})$  and a homomorphism  $\varphi'_\alpha = \left( \varphi \upharpoonright B_{C_* \cup \Lambda_{\eta_\alpha}^{\bar{\xi}_*}} \right) \tau_\alpha \oplus \psi_\alpha \sigma_\alpha^{-1}$ . Let  $\tau' = \sigma_\alpha \sigma^{-1}$ . Notice that

$$\Omega_{\alpha*}^1 \tau' = (\Omega_{\alpha*}^1 \sigma_\alpha) \sigma^{-1} = (\Omega_{\alpha*}^1 \tau_\alpha^{-1}) \sigma^{-1} = V'_* \sigma^{-1} = \Omega_*^1,$$

$$\Omega_{\alpha*}^{2,n} \tau' = \Omega_{\alpha*}^{2,n} \sigma_\alpha \sigma^{-1} = P_{\alpha n} \sigma^{-1} = \Omega_*^{\eta^\alpha \upharpoonright n},$$

so  $Y''_{\alpha*} \tau' = Y''_*$  and  $\tau'$  is  $Y''_{\alpha*}$ -admissible. Furthermore, because of

$$Y'_\alpha \tau' = Y'_\alpha \sigma_\alpha \sigma^{-1} = V' \sigma^{-1} = Y'$$

and

$$(Z_{\alpha,n} \setminus Y'_\alpha) \tau' = Z^{\eta^\alpha \upharpoonright n} \setminus Y',$$

we obtain  $Y''_\alpha \tau' = Y''$ . Similarly,  $X''_\alpha \tau' = X''_*$ , which means that  $(Y''_{\alpha*}, Y''_\alpha, X''_{\alpha*}) \tau' = (Y''_*, Y'', X''_*)$ .

It also follows  $\mathfrak{F}_{Y''_{\alpha*} Y''_\alpha \tau'} = \mathfrak{F}_{Y''_* Y''}$  and

$$\begin{aligned} \varphi'_\alpha \tau' &= \left( \varphi \upharpoonright B_{C_* \cup \Lambda_{\eta_\alpha}^{\bar{\xi}_*}} \right) \tau_\alpha \tau' \oplus \psi_\alpha \sigma_\alpha^{-1} \tau' \\ &= \left( \varphi \upharpoonright B_{C_* \cup \Lambda_{\eta_\alpha}^{\bar{\xi}_*}} \right) \tau \oplus \psi_\alpha \sigma^{-1} \\ &= \left( \varphi \upharpoonright B_{C_* \cup \Lambda_{\eta_\alpha}^{\bar{\xi}_*}} \right) \tau \oplus (\psi \upharpoonright G_\alpha) \\ &= \varphi' \subseteq \psi. \end{aligned}$$

The existence of the  $f$ -closed triples  $(Z''_*, Z'', Y''_*)$  and  $(Y''_*, Y'', X''_*)$  and the fact that  $\psi : G_{J_*J} \rightarrow G_{Z''_*Z''X''_*}$  extends the homomorphism  $\varphi'_\alpha \tau' = \varphi'$  contradicts the previous choice of the elements  $\{b_{\bar{\eta}} \mid \bar{\eta} \in \Lambda^{\bar{\xi}^\alpha}\}$ . □

#### 4. The Final Construction

In this last section, we construct  $\aleph_k$ -free  $A$ -modules  $G$  with a prescribed group endomorphism ring  $\text{End } G = A \oplus \text{Fin } G$ . We not only make use of all our previous work, but we also introduce the Strong Black Box principle and a result involving inessential endomorphisms that is based on ideas from Dugas, Göbel [5].

DEFINITION 61.  $\varphi \in \text{End } G_{Y*Y}$  is *inessential* if  $\overline{B}_{Y*}\varphi \subseteq G_{Y*Y}$ . We denote the set of inessential endomorphisms of  $G_{Y*Y}$  by  $\text{Ines } G_{Y*Y}$ .

LEMMA 62. Let  $\mathcal{V} \subseteq \Lambda$ ,  $\mathfrak{F}_{\Lambda*\mathcal{V}}$  be a regressive family of branch-like elements and  $G = G_{\Lambda*\mathcal{V}}$ . If  $\varphi \in \text{End } G \setminus (A \oplus \text{Ines } G)$ , then there exist a countably-generated  $A$ -submodule  $P$  of  $B$  and some  $z \in \overline{P}$  such that  $z\varphi \notin \langle G, Az \rangle_*$ .

PROOF. Let us begin with the following observation: if  $g \in G$  has infinite support and if  $\overline{\eta} \in [g]_\Lambda$  is of maximal norm, then  $\overline{\eta} \in \Lambda([g])$ . In particular,  $\Lambda([g]) \neq \emptyset$ .

Let  $\mathcal{D} = \{(n, a) \mid n = 0 \text{ or } a \notin p^n A\} \subseteq \omega \times A$ . Our first goal is to find a countably-generated  $A$ -submodule  $P$  of  $B$  such that  $\overline{P}(p^n \varphi - a) \not\subseteq G$  for all  $(n, a) \in \mathcal{D}$ . Let us start by finding a countably-generated  $A$ -submodule  $P'$  of  $B$  such that  $\overline{P}'(\varphi - a) \not\subseteq G$  for all  $a \in A$ , i.e. for all  $(0, a) \in \mathcal{D}$ . Choose some  $w \in \overline{B} \setminus B$  with  $\Lambda([w]) = \emptyset$ ; and let  $P_0 = B_{[w]}$ . Suppose that  $P' = P_0$  fails to have the desired property. Then,  $\overline{P}_0(\varphi - a) \subseteq G$  for some  $a \in A$ . Since  $\varphi - a \notin \text{Ines } G$ , there exists some  $w' \in \overline{B}$  such that  $w'(\varphi - a) \notin G$ . Let  $P_1 = B_{[w] \cup [w']}$ . Towards a contradiction, suppose again that  $\overline{P}_1(\varphi - a') \subseteq G$  for some  $a' \in A$ . Since  $w \in \overline{P}_0 \subseteq \overline{P}_1$ , we have  $w(\varphi - a) - w(\varphi - a') =$

$w(a' - a) \in G$  which is impossible for  $a \neq a'$  because  $\Lambda([w]) = \emptyset$ , so  $w(a' - a)$  cannot be an element of  $G$ . This forces  $a = a'$ , from which we get the desired contradiction

$$G \ni w'(\varphi - a') = w'(\varphi - a) \notin G.$$

Hence, it suffices to take  $P' = P_0$  or  $P' = P_1$ .

Our next task is to find a countably-generated submodule  $P$  of  $B$  containing  $P'$  with the required property. Let

$$\mathcal{N} = \{ n < \omega \mid \exists a_n \in A \left( (n, a_n) \in \mathcal{D} \text{ and } \overline{P'}(p^n \varphi - a_n) \subseteq G \right) \}.$$

Observe that  $a_n$  is unique for  $n \in \mathcal{N}$ : if  $(n, a_n), (n, a'_n) \in \mathcal{D}$ , then

$$w(p^n \varphi - a'_n) - w(p^n \varphi - a_n) = w(a_n - a'_n) \in G,$$

and our previous argument implies that  $a_n = a'_n$ .

Consider  $n \in \mathcal{N}$  and put  $\psi = p^n \varphi - a_n$ . We recursively construct a family  $C = \{ \bar{\nu}_s \mid s < \omega \} \subseteq \Lambda_*$  such that  $\Lambda(C) = \emptyset$ ,  $\max_{r < s} \|e_{\bar{\nu}_r} \psi\| < \|e_{\bar{\nu}_s}\|$  and  $C \cap [\bar{\eta}]$  is finite for all  $\bar{\eta} \in \Lambda$ . Let

$$x_n = \sum_{r=0}^{\infty} p^{(n+1)r} e_{\bar{\nu}_r}.$$

Notice that

$$\sum_{r=s+1}^{\infty} p^{(n+1)r} e_{\bar{\nu}_r} = p^{(n+1)s+n} \sum_{r=1}^{\infty} p^{(n+1)r-n} e_{\bar{\nu}_{s+r}},$$

for every  $s < \omega$ ; and

$$\begin{aligned} x_n \psi &= \left( \sum_{r=0}^{s-1} p^{(n+1)r} e_{\bar{\nu}_r} \psi \right) + p^{(n+1)s} e_{\bar{\nu}_s} (p^n \varphi - a_n) + \left( \sum_{r=s+1}^{\infty} p^{(n+1)r} e_{\bar{\nu}_r} \psi \right) \\ &= \left( \sum_{r=0}^{s-1} p^{(n+1)r} e_{\bar{\nu}_r} \psi \right) - p^{(n+1)s} a_n e_{\bar{\nu}_s} + p^{(n+1)s+n} \left( e_{\bar{\nu}_s} \varphi + \sum_{r=1}^{\infty} p^{(n+1)r-n} e_{\bar{\nu}_{s+r}} \psi \right). \end{aligned}$$

It follows that  $C \subseteq [x_n\psi]$  since  $\max_{r < s} \|e_{\bar{v}_r}\psi\| < \|e_{\bar{v}_s}\|$  and  $a_n \notin p^n A$ . Therefore,  $x_n\psi \notin G$ . We do this construction for every  $n \in \mathcal{N}$ . Let  $P = B_{S \cup [P']}$  where  $S = \bigcup_{n \in \mathcal{N}} [x_n]$ . To verify that  $P$  has the required property, suppose  $\bar{P}(p^n\varphi - a) \subseteq G$  for some  $(n, a) \in \mathcal{D}$ . This means that  $n \in \mathcal{N}$  and  $a = a_n$ , but then again

$$G \ni x_n(p^n\varphi - a) = x_n(p^n\varphi - a_n) \notin G$$

yields a contradiction. Hence  $P$  is as required.

Now take some  $x \in \bar{B} \setminus B$  such that  $[x] \cap ([P] \cup [P\varphi]) = \emptyset$  and  $[x] \cap [\bar{\eta}]$  is finite for all  $\bar{\eta} \in \Lambda$ . If  $x\varphi \in \langle G, Ax \rangle_*$ , then

$$(1) \quad x(p^n\varphi - a) \in G$$

for some  $n < \omega$  and  $a \in A$ . Since  $G$  is pure, we can assume  $(n, a) \in \mathcal{D}$ . Take  $x' \in \bar{P}$  such that  $x'(p^n\varphi - a) \notin G$ . If we assume  $(x + x')\varphi \in \langle G, A(x + x') \rangle_*$ , then

$$(2) \quad (x + x')(p^m\varphi - a') \in G$$

for some  $m \in [n, \omega)$  and  $a' \in A$ . Subtracting (1) multiplied by  $p^{m-n}$  from (2), we obtain

$$(x + x')(p^m\varphi - a') - p^{m-n}x(p^n\varphi - a) = x'(p^m\varphi - a') + x(p^{m-n}a - a') \in G.$$

This yields  $p^{m-n}a = a'$  since  $[x] \cap ([P] \cup [P\varphi]) = \emptyset$  and  $[x] \cap [\bar{\eta}]$  is finite for all  $\bar{\eta} \in \Lambda$ . But then,  $x'(p^m\varphi - a') \in G$  which is a contradiction. Hence we can take  $z = x$  or  $z = x + x'$ .  $\square$

For a group  $G$ , define  $\text{Fin } G$  to the set of endomorphisms of  $G$  with finite rank. Because  $A$  is slender,  $\text{Ines } G = \text{Fin } G$  for the kind of groups we have constructed. We

refer the reader to Göbel, Trlifaj [14], Theorem 12.3.37 for a detailed proof.

We now present the Strong Black Box, which is the last prediction principle that will be used in the final construction. See Göbel, Trlifaj [14] for a proof.

DEFINITION 63. A *trap* for the Strong Black Box is a quintuple  $p = (\eta, V_*, V, \mathfrak{F}, \varphi)$  such that

- (1)  $\eta \in {}^{\omega\uparrow}\lambda_k$ ,
- (2)  $V \in [\Lambda]^{\leq \lambda_{k-1}}$  and  $V_* \in [\Lambda_*]^{\leq \lambda_{k-1}}$ ,
- (3)  $(V_*, V)$  is  $\Lambda$ -closed,
- (4)  $\Lambda^{\eta*} \subseteq V_*$ ,
- (5)  $\|\bar{\xi}\| < \|\eta\|$  for all  $\bar{\xi} \in V \cup V_*$ ,
- (6) For  $\bar{\eta} \in \Lambda$ , if  $\|\bar{\eta}\| < \|\eta\|$  and  $k \notin u_{\bar{\eta}}(V_*)$ , then  $\bar{\eta} \in V$ .
- (7) For  $\bar{\eta} \in \Lambda$ , if  $([\bar{\eta}] \setminus [\bar{\eta} \restriction k]) \cap V_* \neq \emptyset$ , then  $[\bar{\eta}] \subseteq V_*$ .
- (8)  $\mathfrak{F} = \mathfrak{F}_{V_*V} = \{y'_{\bar{\eta}} = b_{\bar{\eta}} + y_{\bar{\eta}} \mid \bar{\eta} \in V, b_{\bar{\eta}} \in \overline{B}_{V_*}\}$  is regressive,
- (9)  $\varphi : G_{V_*V} \rightarrow G_{V_*V}$  is a homomorphism.

THE STRONG BLACK BOX 64. Let  $\mu$  be an infinite cardinal,  $\lambda = \mu^+$ ,  $\theta \leq \lambda$  such that  $\mu^\theta = \mu$  and  $k > 1$ . If  $E \subseteq \lambda^\circ$  is stationary, then there is a family

$$\{p_\alpha = (\eta^\alpha, V_{\alpha*}, V_\alpha, \mathfrak{F}_\alpha, \varphi_\alpha) \mid \alpha < \lambda\}$$

of traps such that

- (1)  $\|\eta^\alpha\| \in E$  for all  $\alpha < \lambda$ ,



- (2)  $\|\eta^\alpha\| \leq \|\eta^\beta\|$  for all  $\alpha < \beta < \lambda$ ,
- (3) If  $\|\eta^\alpha\| = \|\eta^\beta\|$  for  $\alpha \neq \beta$ , then  $\|V_{\alpha*} \cap V_{\beta*}\| < \|\eta^\alpha\|$ ,
- (4) For any  $\mathcal{V} \subseteq \Lambda$ , any regressive family  $\mathfrak{F}_{\Lambda*\mathcal{V}} = \{y'_{\bar{\eta}} = b_{\bar{\eta}} + y_{\bar{\eta}} \mid \bar{\eta} \in \mathcal{V}, b_{\bar{\eta}} \in \overline{B}\}$ , any  $\varphi \in \text{End } G_{\Lambda*\mathcal{V}}$ ,  $U \in [\Lambda_*]^{\leq \theta}$  and  $\delta < \lambda$ , the set

$$\{\gamma \in E \mid \exists \alpha < \lambda (\|\eta^\alpha\| = \gamma, \delta < 0\eta^\alpha, V_\alpha = \mathcal{V}_{V_{\alpha*}}, \mathfrak{F}_\alpha = \mathfrak{F}_{\Lambda*V_\alpha}, \varphi_\alpha \subseteq \varphi, U \subseteq V_{\alpha*})\}$$

is unbounded, where  $V_\alpha = \mathcal{V}_{V_{\alpha*}}$  is only possible for  $k > 1$ .

MAIN THEOREM 65. If  $A$  is a ring with free additive group  $A^+ = \bigoplus_{\alpha < \kappa} \mathbb{Z}e_\alpha$  such that  $\overline{A} = \widehat{A} \cap \prod_{\alpha < \kappa} \mathbb{Z}e_\alpha$  is an  $A$ -module,  $|A| < \mu$ ,  $k \in (0, \omega)$  and  $\lambda = \beth_k^+(\mu)$ , then it is possible to construct an  $\aleph_k$ -free  $A$ -module  $G$  of cardinality  $\lambda$  such that  $G$  is separable as an abelian group and  $\text{End } G = A \oplus \text{Ines } G = A \oplus \text{Fin } G$ .

PROOF. Since the case  $k = 1$  is a classical result due to M. Dugas and R. Göbel (see [5]), we assume  $k > 1$ . Consider the stationary subset  $\lambda_k^o = \{\alpha < \lambda_k \mid \text{cf}(\alpha) = \omega\}$  of  $\lambda_k = \mu_k^+$ . By Solovay's Theorem we can decompose  $\lambda_k^o$  into  $\lambda_k$  disjoint stationary subsets, say  $\lambda_k^o = \dot{\bigcup}_{\alpha < \lambda_k} E_\alpha$ . Now let

$$\mathcal{I} = \{(P, z) \mid P \leq B \text{ is countably-generated}, z \in \overline{P} \setminus P\},$$

and notice that  $|\mathcal{I}| = \lambda_k$ . Therefore, we are allowed to write

$$\lambda_k^o = \dot{\bigcup}_{(P, z) \in \mathcal{I}} E_{(P, z)}.$$

For each  $E_{(P, z)}$ , the Strong Black Box provides us with a family of traps

$$p_\alpha^{(P, z)} = (\eta^{\alpha(P, z)}, V_{\alpha*}^{(P, z)}, V_\alpha^{(P, z)}, \mathfrak{F}_\alpha^{(P, z)}, \varphi_\alpha^{(P, z)})$$

for  $\alpha < \lambda_k$ . We gather all these traps and order them according to the norm of their first component, namely  $\|\eta^\alpha\| \leq \|\eta^\beta\|$  for all  $\alpha < \beta < \lambda_k$ .

Let  $\mathcal{V} = \bigcup_{\alpha < \lambda_k} \Lambda^{\eta^\alpha}$  (see Definition 56). We will construct a regressive family  $\mathfrak{F}_{\Lambda^* \mathcal{V}}$  of branch-like elements by choosing for all  $\alpha < \lambda_k$  and all  $\bar{\eta} \in \Lambda^{\eta^\alpha}$  an element  $b_{\bar{\eta}} \in \overline{B}$ , and define  $G = G_{\Lambda^* \mathcal{V}}$ . If we get, when considering the trap  $p_\alpha = (\eta^\alpha, V_{\alpha^*}, V_\alpha, \mathfrak{F}_\alpha, \varphi_\alpha)$  and the unique  $(P, z) \in \mathcal{I}$  such that  $\|\eta^\alpha\| \in E_{(P, z)}$ , that  $[P] \subseteq V_{\alpha^*}$ ,  $\|z\| < 0\eta^\alpha$  and  $z\varphi_\alpha \notin G_{V_{\alpha^*}V_\alpha}$ , we will obtain the  $b_{\bar{\eta}}$ 's for  $\bar{\eta} \in \Lambda^{\eta^\alpha}$  by means of the Step Lemma 60. Otherwise we will set  $b_{\bar{\eta}} = 0$  for all  $\bar{\eta} \in \Lambda^{\eta^\alpha}$ .

Let  $\alpha < \lambda_k$  and  $(P, z)$  be the unique element of  $\mathcal{I}$  such that  $\|\eta^\alpha\| \in E_{(P, z)}$ . Assume  $[P] \subseteq V_{\alpha^*}$ ,  $\|z\| < 0\eta^\alpha$  and  $z\varphi_\alpha \notin G_{V_{\alpha^*}V_\alpha}$ . Choose two elements  $z_1, z_2 \in P$  with infinite support such that  $z_1 + z_2 = z$ ,  $[z_1] \dot{\cup} [z_2] = [z]$  and  $u_{\bar{\eta}}([z_1]) = u_{\bar{\eta}}([z_2]) = [1, k]$  for all  $\bar{\eta} \in \Lambda$ . Without loss of generality,  $z_1\varphi_\alpha \notin G_{V_{\alpha^*}V_\alpha}$ . Let  $V_* = V_{\alpha^*} \dot{\cup} \Lambda_*^{\eta^\alpha}$ ,  $V = V_\alpha \dot{\cup} \Lambda^{\eta^\alpha}$  and  $U_* = [\Lambda^{\eta^\alpha}] \cup [z_1]$ . These unions are indeed disjoint because  $V_{\alpha^*} \cap \Lambda_*^{\eta^\alpha} = V_\alpha \cap \Lambda^{\eta^\alpha} = \emptyset$  by Definition 63(5).

We claim that  $(V_*, V, U_*)$  is  $k$ -closed: Since  $(V_{\alpha^*}, V_\alpha)$  is  $\Lambda$ -closed and  $\Lambda^{\eta^\alpha} \subseteq V_{\alpha^*}$ , it follows that  $(V_*, V)$  is  $\Lambda$ -closed. If  $\bar{\eta} \in V$ , then either  $\bar{\eta} \in V_\alpha$  or  $\bar{\eta} \in \Lambda^{\eta^\alpha}$ . If  $\bar{\eta} \in V_\alpha$ , then in particular  $\eta_k \neq \eta^\alpha$  since  $\|\bar{\eta}\| < \|\eta^\alpha\|$ , so  $[\bar{\eta}]_n \cap [\Lambda^{\eta^\alpha}] = \emptyset$  for all large enough  $n < \omega$ . Moreover, for all  $m \in [1, k]$ ,  $[\bar{\eta} \upharpoonright m]_n \not\subseteq [z_1]$  since  $u_{\bar{\eta}}([z_1]) = [1, k]$ . Hence,  $u_{\bar{\eta}}(U_*) = [1, k]$  for all  $\bar{\eta} \in V_\alpha$ . If  $\bar{\eta} \in \Lambda^{\eta^\alpha}$ , then  $[\bar{\eta}] \subseteq [\Lambda^{\eta^\alpha}]$  and  $\bar{\eta} \in V_{U_*}$ . This proves our claim. Moreover,  $\Lambda^{\eta^\alpha} = V_{U_*}$ .

Let  $\mathfrak{F}_{V_* V} = \mathfrak{F}_\alpha \cup \mathfrak{F}_{V_* V_{U_*}}$  where  $\mathfrak{F}_{V_* V_{U_*}} = \{\varepsilon_{\bar{\eta}} z_1 + y_{\bar{\eta}} \mid \bar{\eta} \in \Lambda^{\eta^\alpha}\}$  for some  $\varepsilon_{\bar{\eta}} \in \{0, 1\}$ . Observe that if  $\bar{\eta} \in V_{U_*}$ , then either  $b_{\bar{\eta}} = 0$  or  $[b_{\bar{\eta}}] = [z_1] \subseteq U_*$ . Hence,  $\mathfrak{F}_{V_* V}$  is

$(V, U_*)$ -suitable. Put  $f = k - 1$ ,  $\bar{\xi} = \eta^\alpha$ ,  $C_* = [z_1]$  and

$$\psi_\alpha = (\varphi_\alpha \upharpoonright B_{I_*(\bar{\xi})})\rho_{V_*U_*} : B_{I_*(\bar{\xi})} \rightarrow G_{V_*VU_*}$$

(recall that  $(J_*(\bar{\xi}), J(\bar{\xi}), I_*(\bar{\xi})) = (C_* \cup [\Lambda^{\bar{\xi}}], \Lambda^{\bar{\xi}}, C_* \cup \Lambda^{\bar{\xi}*})$  as in the Step Lemma 60), and notice  $z_1\psi_\alpha \notin G_{V_*VU_*}$  since  $z_1\varphi_\alpha \notin G_{V_{\alpha*}V_\alpha}$ . We now choose the elements  $b_{\bar{\eta}} \in \bar{B}_{J_*}$  for all  $\bar{\eta} \in J = \Lambda^{\eta^\alpha}$  by applying the Step Lemma 60 (remember that the choice of these elements is independent from  $\mathfrak{F}_{V_*VU_*}$ ).

To obtain a contradiction, suppose that there is some  $\varphi \in \text{End } G \setminus (A \oplus \text{Ines } G)$ . We apply Lemma 62 to  $G$  to obtain a countably-generated submodule  $P$  of  $B$  and some  $z \in \bar{P}$  such that  $z\varphi \notin \langle G, Az \rangle_*$ . Then,  $z \in \bar{P} \setminus P$ ; and  $(P, z)$  is an element of the index set  $\mathcal{I}$ . Since the set

$$\{ \gamma \in E_{(P,z)} \mid \exists \alpha < \lambda_k ( \| \eta^{\alpha(P,z)} \| = \gamma, \| z \| < 0\eta^{\alpha(P,z)}, V_\alpha^{(P,z)} = \mathcal{V}_{V_{\alpha*}^{(P,z)}},$$

$$\mathfrak{F}_\alpha^{(P,z)} = \mathfrak{F}_{\Lambda_*V_\alpha^{(P,z)}}, \varphi_\alpha^{(P,z)} \subseteq \varphi, [P] \subseteq V_{\alpha*}^{(P,z)} \}$$

is unbounded by the Strong Black Box, we can choose an ordinal  $\gamma$  from this set with the corresponding trap  $p_{\alpha'}^{(P,z)}$ . Pick  $\alpha < \lambda_k$  such that  $p_{\alpha'}^{(P,z)} = p_\alpha$ . Because of  $[P] \subseteq V_{\alpha*}$ ,  $\| z \| < 0\eta^\alpha$  and  $z\varphi_\alpha \notin G_{V_{\alpha*}V_\alpha} \subseteq G$ , the conditions for applying the Step Lemma 60 are satisfied. Take  $z_1$  and  $z_2$  as before. Let  $(Z_*, Z) = (\Lambda_*, \mathcal{V})$ , which is clearly  $\Lambda$ -closed, and  $(Y_*, Y, X_*) = (V_*, V, U_*)$ . We have already seen that  $(Y_*, Y, X_*)$  is  $k$ -closed. To prove that  $(Z_*, Z, Y_*)$  is also  $k$ -closed, let  $\bar{\eta} \in Z = \bigcup_{\alpha < \lambda_k} \Lambda^{\eta^\alpha}$ . Thus,  $\eta_k = \eta^\beta$  for some  $\beta < \lambda_k$ . If  $\| \eta^\beta \| > \| \eta^\alpha \|$ , then  $u_{\bar{\eta}}(Y_*) = [1, k]$  since  $\| Y_* \| = \| \eta^\alpha \|$ . Now assume  $\| \eta^\beta \| < \| \eta^\alpha \|$  and  $|u_{\bar{\eta}}(Y_*)| < k$ . Notice that, in this case,  $u_{\bar{\eta}}(Y_*) = u_{\bar{\eta}}(V_{\alpha*})$ . If

$k \notin u_{\bar{\eta}}(Y_*)$ , then  $\bar{\eta} \in V_\alpha = (\mathcal{V})_{V_{\alpha*}} \subseteq (\mathcal{V})_{Y_*}$  by Definition 63(6) and the  $\Lambda$ -closedness of  $(V_{\alpha*}, V_\alpha)$ . Hence,  $\bar{\eta} \in Z_{Y_*}$ . If  $m \notin u_{\bar{\eta}}(Y_*)$  for some  $m \neq k$ , then  $\bar{\eta} \upharpoonright \langle m, n \rangle \in V_{\alpha*}$  for some  $n < \omega$ , which implies that  $[\bar{\eta}] \subseteq V_{\alpha*}$  by Definition 63(7). Consequently,  $\bar{\eta} \in Z_{Y_*}$  as well. Finally, assume  $\|\eta^\beta\| = \|\eta^\alpha\|$ . If  $\beta = \alpha$ , then  $[\bar{\eta}] \subseteq Y_*$ , so  $\bar{\eta} \in Z_{Y_*}$ . If  $\beta \neq \alpha$ , then  $\bar{\eta} \upharpoonright \langle m, n \rangle \notin Y_*$  for all  $m \neq k$  and  $n < \omega$  since  $\|\bar{\eta} \upharpoonright \langle m, n \rangle\| = \|\eta^\alpha\|$ . This implies  $[1, k-1] \subseteq u_{\bar{\eta}}(Y_*)$ . If  $k \notin u_{\bar{\eta}}(Y_*)$ , then  $[\bar{\eta} \upharpoonright k]_n \subseteq V_{\alpha*}$  for some  $n < \omega$ . By Definition 63(4), we also have  $[\bar{\eta} \upharpoonright k]_n \subseteq V_{\beta*}$ . This yields  $[\bar{\eta} \upharpoonright k]_n \subseteq V_{\alpha*} \cap V_{\beta*}$  and  $\|V_{\alpha*} \cap V_{\beta*}\| = \|\eta^\alpha\|$  contradicting the Strong Black Box 64. Therefore,  $u_{\bar{\eta}}(Y_*) = [1, k]$ . This proves that  $(Z_*, Z, Y_*)$  is  $k$ -closed.

Now we verify  $Z_{Y_*} = Y$ . It is immediate that  $Y \subseteq Z_{Y_*}$ . Let  $\bar{\eta} \in Z_{Y_*}$  with  $\eta_k = \eta^\beta$  for some  $\beta < \lambda_k$ . If  $\beta = \alpha$ , then  $\bar{\eta} \in \Lambda^{\eta^\alpha} \subseteq Y$ . If  $\beta \neq \alpha$ , then  $[\bar{\eta}]_n \subseteq V_{\alpha*}$  for some  $n < \omega$ . In particular  $\|\bar{\eta}\| < \|\eta^\alpha\|$  and  $u_{\bar{\eta}}(V_{\alpha*}) = \emptyset$ . By Definition 63(6),  $\bar{\eta} \in V_\alpha$ , so  $\bar{\eta} \in Y$  and  $Z_{Y_*} \subseteq Y$ .

Finally, we check that  $\mathfrak{F}_{\Lambda_*\mathcal{V}}$  is  $(Z, Y_*)$ -suitable and  $\mathfrak{F}_{\Lambda_*Y}$  is  $(Y, X_*)$ -suitable. If  $\bar{\eta} \in Z_{Y_*} = Y$ , then  $\bar{\eta} \in V_\alpha$  or  $\bar{\eta} \in \Lambda^{\eta^\alpha}$ . In the first case,  $b_{\bar{\eta}} \in \overline{B}_{V_{\alpha*}}$  by the definition of  $\mathfrak{F}_\alpha$ . Thus,  $[b_{\bar{\eta}}] \subseteq V_{\alpha*}$ . In the second case,  $b_{\bar{\eta}} = z_1$ ; and  $[b_{\bar{\eta}}] \subseteq X_* \subseteq Y_*$  follows. This proves that  $\mathfrak{F}_{\Lambda_*\mathcal{V}}$  is  $(Z, Y_*)$ -suitable. Moreover, the second case also yields that  $\mathfrak{F}_{\Lambda_*Y}$  is  $(Y, X_*)$ -suitable since  $Y_{X_*} = V_{U_*} = \Lambda^{\eta^\alpha}$ . Hence, the homomorphism  $\varphi\rho_{Z_*X_*} = \varphi\rho_{\Lambda_*U_*}$  extends the homomorphism

$$\psi_\alpha = (\varphi_\alpha \upharpoonright B_{I_*(\bar{\xi})})\rho_{Y_*X_*} = (\varphi_\alpha \upharpoonright B_{I_*(\eta^\alpha)})\rho_{V_*U_*}.$$

The existence of  $(Z_*, Z, Y_*)$ ,  $(Y_*, Y, X_*)$ ,  $\mathfrak{F}_{\Lambda_*\mathcal{V}}$ ,  $\tau = \text{Id}$  and  $\psi_\alpha \subseteq \varphi\rho_{Z_*X_*}$  contradicts the choice of the  $b_{\bar{\eta}}$ 's for  $f = k-1$ ,  $\bar{\xi} = \eta^\alpha$ ,  $C_* = [z_1]$  and  $z_1$ .  $\square$

## References

- [1] A.L.S. Corner, *Every countable reduced torsion-free ring is an endomorphism ring*, Proc. London Math. Soc. (3) **13** (1963), 687 - 710.
- [2] A.L.S. Corner, R. Göbel, *Prescribing endomorphism algebras - A unified treatment*, Proc. London Math. Soc. (3) **50** (1985), 447 - 479.
- [3] A.L.S. Corner, R. Göbel, *Subgroups of the Baer-Specker group with prescribed endomorphism ring and large dual*, Proceeding in honour of Adalberto Orsatti's 60th birthday, Abelian groups, module theory, and topology, ed. D. Dikranjan, L. Salce, Marcel Dekker, New York (1998), 113 - 123.
- [4] M. Dugas, R. Göbel, *Every cotorsion-free ring is an endomorphism ring*, Proc. London Math. Soc. (3) **45** (1982), 319 - 336.
- [5] M. Dugas, R. Göbel, *Endomorphism rings of separable torsion-free abelian groups*, Houston J. Math. **11** (1985), 471 - 483.
- [6] D.S. Dummit, R.M. Foote, *Abstract Algebra* Third ed., John Wiley & Sons Inc. (2004).
- [7] P. C. Eklof, A. H. Mekler, *Almost Free Modules*, Revised ed., North-Holland, New York (2002).
- [8] R. Engelking, *General Topology*, Revised and completed ed., Sigma Series in Pure Mathematics **6**, Heldermann Verlag, Berlin (1989).
- [9] P. C. Eklof, S. Shelah, *The Kaplansky test problems for  $\aleph_1$ -separable groups*, Proc. Amer. Math. Soc. **126** (1998), 1901 - 1907.
- [10] L. Fuchs, *Infinite Abelian Groups - Vol. 1 & 2*, Academic Press, New York (1970, 1973).
- [11] R. Göbel, D. Herden, S. Shelah, *Prescribing endomorphism algebras of  $\aleph_n$ -free modules*, submitted (2010).
- [12] R. Göbel, S. Shelah,  *$\aleph_n$ -free modules with trivial duals*, Result. Math. **54** (2009), 53 - 64.

- [13] R. Göbel, S. Shelah, L. Strüngmann,  $\aleph_n$ -free modules over complete discrete valuation domains with small dual, to appear in Glasgow Journal of Mathematics.
- [14] R. Göbel, J. Trlifaj, *Approximations and Endomorphism Algebras of Modules*, Expositions in Mathematics **41**, Walter de Gruyter Verlag, Berlin (2006).
- [15] K. R. Goodearl, P. Menal, J. Moncasi, *Free and residually Artinian regular rings*, J. Algebra **156** (1993), 407 - 432.
- [16] P.A. Griffith,  $\aleph_n$ -free abelian groups, Quart. J. Math. (Oxford) (2) **23** (1972), 417 - 425.
- [17] P. Hill, *New criteria for freeness in abelian groups II*, Trans. Amer. Math. Soc. **196** (1974), 191 - 201.
- [18] T. Jech, *Set Theory*, The Third Millennium Edition, revised and expanded, Springer Monographs in Mathematics, Springer, Berlin Heidelberg (2002).
- [19] P. P. Nielsen, *Row and Column finite matrices*, Proc. Amer. Math. Soc. **135** (2007), 2689 - 2697.
- [20] S. Shelah,  $\aleph_n$ -free abelian groups with no non-zero homomorphisms to  $\mathbb{Z}$ , Cubo - A Mathematical Journal **9** (2007), 59 - 79.